

FIRST DRAFT Released for Public Comment on July 3, 2019



Striving for Zero

CALIFORNIA'S STRATEGIC PLAN FOR SUICIDE PREVENTION 2020 – 2025

Support for people at risk for suicide or those supporting people at risk is available by calling the National Suicide Prevention Lifeline 1-800-273-TALK (8255)

About the Commission

The Mental Health Services Oversight and Accountability Commission, an independent state agency, was created in 2004 by voter-approved Proposition 63, the Mental Health Services Act. California voters created the Commission to provide oversight, accountability, and leadership to guide the transformation California's mental health system. The 16-member Commission is composed of one Senator, one Assembly member, the State Attorney General, the State Superintendent of Public Instruction, and 12 public members appointed by the Governor. By law, the Governor's appointees are people who represent different sectors of society, including individuals with mental health needs, family members of people with mental health needs, law enforcement, education, labor, business, and the mental health profession.

Commissioners

KHATERA TAMPLEN, Chair; ***Suicide Prevention Subcommittee Member***; Consumer Empowerment Manager, Alameda County Behavioral Health Care Services

LYNNE ASHBECK, Vice Chair; Senior Vice President of Community Engagement and Population Wellness, Valley Children's Healthcare

MAYRA E. ALVAREZ; President, Children's Partnership

RENEETA ANTHONY; Executive Director, A3 Concepts LLC

JIM BEALL; California State Senator, District 15

KEN BERRICK; Chief Executive Officer, Seneca Family of Agencies

JOHN BOYD, Psy.D.; Chief Executive Officer of Mental Health Services, Sutter Health Care

BILL BROWN; Sheriff, County of Santa Barbara

KEYONDRIA BUNCH, Ph.D.; Clinical Psychologist, Emergency Outreach Bureau, Los Angeles County Department of Mental Health

WENDY CARRILLO; California State Assemblymember, District 51

ITAI DANOVITCH, M.D.; Chair, Department of Psychiatry and Behavioral Neurosciences, Cedars-Sinai Medical Center

DAVID GORDON; Superintendent, Sacramento County Office of Education

MARA MADRIGAL-WEISS; ***Suicide Prevention Subcommittee Member***; Director of Wellness and Student Achievement, Student Services and Programs Division, San Diego County Office of Education

GLADYS MITCHELL; Former Staff Services Manager, California Department of Health Care Services and California Department of Alcohol and Drug Programs

TINA WOOTON; ***Suicide Prevention Subcommittee Chair***; Consumer Empowerment Manager, Santa Barbara County Department of Behavioral Wellness

TOBY EWING, Ph.D.; Executive Director

BRIAN SALA, Ph.D.; Deputy Director of Evaluation and Program Operations

DAWNTÉ EARLY, Ph.D.; Chief of Research and Evaluation

ASHLEY MILLS, M.S.; Senior Researcher and Suicide Prevention Project Lead

Acknowledgment

This plan is dedicated to people lost to suicide and people living with suicidal thoughts and actions, and their loved ones. The Commission would like to express its thanks to the many survivors, community members, family members, administrators, providers, researchers, and policymakers who contributed to the development of this plan. We greatly appreciate the time, commitment, and energy devoted to exploring the challenges and solutions surrounding efforts to prevent suicide.

We would like to extend a special thank you to the survivors of suicide attempt and loss who bravely and honestly shared their stories, experiences, and unique insights into opportunities to improve suicide prevention strategies. Many people are affected by suicide, including Commissioners and staff directly involved in the development of this plan. The Commission affirms the urgency of putting in place sound strategies to prevent further loss of life.

Lives can be saved. There is hope.

Get Help Now

If you or someone else needs support, a trained crisis counselor can be reached by calling the National Suicide Prevention Lifeline at **800-273-TALK (8255)** or by texting TALK to **741741**.

- For veterans, call the Lifeline at **800-273-TALK (8255)** and press 1.
- For LGBTQ people, call The Trevor Project at **866-488-7386** or text START to **678678**.
- For transpeople, call the Trans Lifeline at **877-565-8860**.
- For people who are deaf or hard of hearing, call the Lifeline at **800-799-4889**.
- For law enforcement personnel, call the COPLINE at **800-267-5463**.

All of the above are confidential resources, available 24 hours a day, seven days a week.

Assessing for suicide risk uses a collaborative and transparent approach to providing support services and referral options.

If someone is showing warning signs or communicating a desire to die, take the following steps:

1. **ASK** “Are you thinking about suicide or feeling that life may not be worth living?” and assess the person’s safety by asking if the person has a specific plan and any intent to act on that plan. Ask if the person has already begun acting on these thoughts or made a suicide attempt. Risk of death by suicide increases significantly as people put more pieces of a plan in place.
2. **EXPRESS** care. Suicidal desire may be a frightening and isolating experience. Express care to emphasize that help is available, including confidential resources.

3. **REACH OUT** for support by calling the crisis lines (see above) to be connected to resources. All crisis lines are available for people in crisis OR those supporting people in crisis.
4. **FOLLOW-UP** by calling, texting, or visiting to ask how the person is doing and if additional support is needed.

For more information or resources visit www.SPRC.org and www.suicideispreventable.org.

Plan Contents

Executive Summary	6
Stigma and Myths	10
Introduction	11
Strategic Aims, Goals, and Action	13
Strategic Aim 1: Establish infrastructure to support suicide prevention strategies	15
Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness	25
Strategic Aim 3: Enhance early identification of suicide risk and increase access to services based on risk	37
Strategic Aim 4: Advance suicide-specific services and supports	45
Plan Development	51
Suicidal Behavior: Definitions, Theory, and Key Concepts for Prevention	60
Assessing for Suicide Risk	62
Suicide Theory	62
Means Matter	63
Inherent Challenges and Emerging Innovations	64
Suicidal Behavior in California	67
Suicide Data	67
Other Suicidal Behavior Data	69
Data Limitations	69
Risk and Protective Factors	71
Cultural Considerations	71
Vulnerable Groups	74
Best Practice in Suicide Prevention	78
Universal Prevention Strategies	78
Selective Prevention Strategies	84
Indicated Prevention Strategies	87
References	91

Executive Summary

More than 47,000 Americans lose their lives to suicide each year. While global suicide rates are decreasing, the national suicide rate has been on a steady rise since 1999. Suicide is the tenth leading cause of death in the United States, but ranks as the second leading cause for people ages 10 to 34. Each year an estimated 25 suicide attempts occur for every death by suicide; among youth, up to 200 attempts occur for every suicide death. In 2017, the national suicide rate was 14 per 100,000 people. California's suicide rate – 10.7 per 100,000 residents – is lower relative to other states, but includes much higher rates in certain counties and demographic groups.

Suicide is a complex public health challenge involving many biological, psychological, social, and cultural determinants. Major risk factors for suicide are prior suicide attempt; substance use disorder; mood disorders, such as depression; and access to methods to attempt suicide. Common factors that lessen risk for suicide are access to effective medical and mental health care; connectedness to others; problem-solving skills; and contacts, such as postcards or letters, from service providers and caregivers. While women and youth of color *attempt* suicide at greater rates relative to other groups, middle-aged and older white men *die* by suicide at greater rates. In the U.S., nearly 7 out of 10 suicides are by white men. The most common method for suicide attempt is drug overdose, while firearms are the most common means for suicide death.

Misconceptions about suicidal behavior continue to challenge prevention efforts, despite advancements in the study of suicide and its prevention. These misconceptions include pervasive myths that may prevent people at risk from seeking help and discourage people from asking loved ones about suicide risk. Internal suffering that accompanies the desire to die may remain hidden unless a person is directly asked about their needs. Misconceptions continue as strategies to reduce a person's access to potentially lethal methods of injury are common in other prevention fields, yet they remain underutilized in suicide prevention. Physical barriers on bridges, locking doors on railways, and locking windows positioned at lethal heights prevents accidental *and* intentional falling resulting in injury or death. Likewise, safely storing guns in the home prevents accidental *and* intentional injury and death among children and adults.

Prevention efforts must address dynamic risk factors, which can change over a person's lifetime. Research on the variability in risk and protective factors among vulnerable groups is underway, but much remains unknown. In addition, suicide prevention requires engagement of private and public partners across multidisciplinary fields, posing unique challenges to wide-scale collaborations for integrated planning. Efforts are further complicated by inconsistent definitions of suicidal behavior, which affect data monitoring. Lastly, assessing for risk is not currently uniform, challenging suicide risk detection, which is constrained by significant ethical, training, and legal considerations.

Suicidal Behavior in California, 2017

- 4,323 people died by suicide
- 108,075 estimated suicide attempts
- 18,153 people visited or were admitted to the emergency department for intentional self-harm
- Over 1.1 million adults reported serious thoughts of suicide

Notwithstanding the challenges to prevention, research demonstrates that lives can be saved using effective interventions, and that public health strategies can prevent loss of life on a broad scale. Suicide prevention efforts must be centered on the interacting and repeating cycle of four core elements in the Public Health Model: 1) using data to define the problem, 2) identifying factors that increase and reduce risk, 3) developing interventions and testing their effectiveness, and 4) scaling-up effective interventions through continued evaluation and broad dissemination.

In early 2018, California's Mental Health Services Oversight and Accountability Commission launched an effort to develop a suicide prevention plan for the State of California. The last plan was developed in 2008. Under the leadership of a subcommittee chaired by Commissioner Tina Wooton, the Commission engaged national and local experts, reviewed research, and convened public hearings and forums, where community members, policy leaders, and those with lived experience provided guidance and insight. The goal was to develop an achievable policy agenda and a foundation for suicide prevention based on best practices. The Commission's objective is to equip and empower California communities with the information they need to minimize risk, improve access to care, and prevent suicidal behaviors.

While the state can support local communities and assume a leadership role, the success of any strategic plan depends on the integrated efforts of private and public partners. This synergy is already taking place on many fronts. Private and public health care systems are moving toward integration with behavioral health systems and providers. Public health leaders are investigating risk factors for suicide and novel interventions for its prevention, within communities and service delivery systems. Schools are working with local leaders to increase access to mental health services and deliver social emotional learning that will benefit a student over a lifetime. Businesses are recognizing the importance of workplace wellbeing and expanding pathways to support through modern employee assistance programs.

Fueled by this synergy, California's suicide prevention plan is framed by four strategic aims:

STRATEGIC AIM 1: Establish infrastructure to support suicide prevention strategies

Infrastructure includes visible leadership and networked partnerships, effective management of assets and resources, and data monitoring and evaluation. Leadership is particularly vital to

establish suicide prevention as a priority public health outcome, and to provide oversight to hold systems accountable. Resources must be integrated and coordinated with clearly defined roles and responsibilities among partners, while data must be collected, monitored, and used to identify what works to promote continuous quality improvement.

STRATEGIC AIM 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness Examples include creating safe environments by erecting suicide deterrent systems at bridges where suicides are known to occur, teaching resiliency skills in early education to prepare youth to manage stressors, and creating peer groups to reduce stigma and isolation and increase the visibility of available services and supports. Media, including the entertainment industry, can prevent suicide through responsible reporting of suicide death, destigmatizing mental illness, and by highlighting mental health resources.

STRATEGIC AIM 3: Increase early identification of suicide risk and access to services based on risk Trainings to help people recognize the warning signs of suicide and to have safe conversations about suicide with people at risk are available and have demonstrated effectiveness. People trained to detect suicide risk and safely intervene can be integrated into settings where members of vulnerable groups live, work, learn, and receive care. Screening tools can identify people at risk for suicide, while brief interventions – just like those regularly used for problem alcohol use – empower people at risk to recognize their personal warning signs, identify coping strategies and a supportive social network, reduce access to lethal means, and reach out for professional support so that suicidal crises can be managed. Crisis services and support should be widely available, accessible, and varied in order to benefit the diverse range of people in need of help.

STRATEGIC AIM 4: Advance suicide-specific services and supports Providers should be equipped to help those at risk and trained to deliver care that reflects best practices. Low-cost, high-impact post-hospitalization postcards and referral services are effective strategies for preventing future suicidal behavior and should be a standard component of aftercare. Swift response to support families, loved ones, and, in some cases, entire communities, must follow every suicide.

The state should take the following actions to support long-term strategic aims:

- Establish leadership through an Office of Suicide Prevention and oversight through the California Suicide Prevention Council
- Support the development of local suicide prevention strategic planning and implementation
- Centralize timely suicidal behavior data to guide efforts and innovate new practices
- Create safe environments by reducing access to lethal means
- Increase resiliency and seeking services for behavioral health needs

- Advance understanding of how efforts to prevent of other forms of violence and trauma may reduce suicidal behavior
- Promote training on and adherence to guidelines for safe messaging and reporting about suicide in the media and entertainment industry
- Enhance uniform screening for suicide risk across health care settings, and require suicide prevention training for licensed providers in all hospital settings
- Prioritize the delivery of a continuum of crisis services and expand capacity as needed
- Create a certification for behavioral health providers delivering best practice to treat suicide risk
- Require follow-up and continuity of care for people discharged from hospital settings after receiving suicide-related services
- Ensure systematic and respectful response following a suicide loss

Striving for zero – the elimination of suicide in California – will demand leadership, commitment, and honest conversations about suicide risk and resiliency, as well as barriers that disrupt suicide prevention efforts. This plan outlines public health aims aligned with nationally directed strategies and calls for crucial advancements in innovation and health care access to develop and integrate practices capable of helping millions of people. California has the ingenuity, capacity, and leadership to take a decisive stand against suicide. One life lost to suicide is too many, so let's begin now.

Stigma and Myths

Stigma is a major obstacle to preventing suicide. Stigma refers to negative attitudes and beliefs about people with behavioral health needs. Such needs include problem substance use and problem eating, serious psychological distress, and mental illness, and their severity can range from distress to diagnosable illnesses and disorders. Stigma not only discourages people from seeking help, but also can prevent people, families, and communities from becoming connected with meaningful support. Stigma also affects the reporting and recording of suicides and the circumstances leading up to a suicide, such as a previous attempt or death in the family. Consequently, prevention efforts are stymied by the underreporting of suicidal behavior. To demonstrate one tactic that can combat stigma, the Commission uses non-stigmatizing language throughout this plan. Stigmatizing language includes the phrases *committed suicide*, *completed or successfully completed suicide*, *suicidal person*, *unsuccessful or failed suicide attempt*, and *mentally ill*.

Non-Stigmatizing:	Stigmatizing:
Died by suicide	Committed suicide
Person at risk of suicide	Suicidal person
Person living with mental illness	Mentally ill person


Myths and misconceptions about the prevention of suicide may hinder prevention efforts.¹ Below are common examples of these myths and the facts associated with each.

MYTH	FACT
Most suicides are impulsive and happen without warning.	Over 70 percent of people who die by suicide communicated to someone their plans for the attempt prior to death. Planning, including obtaining the means by which to attempt suicide and identifying a location, often happens well before the attempt – sometimes years in advance. ² Most suicides are preceded by warning signs, such as communicating the desire to die, of having no reason to live, or the feeling of being a burden. ³
People who want to die are determined and there is no changing their minds.	Over 90 percent of people who were interrupted in a suicide attempt will not go on to die by suicide at another location or by other methods. ⁴ Research suggests that those at risk for suicide show extreme ambivalence about the desire to die or live, and express a high degree of suffering. Attempt survivor accounts suggest people are relieved to have lived through an attempt and regain their desire to live. ⁵ This fact highlights the opportunity to intervene and separate the person at risk from lethal means for a suicide attempt.
Communicating about suicide will plant the seed for suicidal thoughts, increasing risk.	Communicating openly about suicide and asking about risk has been shown to be lifesaving. It encourages people to seek help, promotes a sense of belonging, and connects people to care.

Introduction

Suicide is a serious public health challenge, accounting for nearly 800,000 deaths each year worldwide.⁶ In the United States, suicide remains among the top ten causes of death, claiming twice as many lives each year as homicide. Suicide rates have remained relatively intractable nationally over the past 50 years, and rose 33 percent between 1999 and 2017 – from 10.5 to 14 per 100,000 Americans.⁷ It is estimated that for every suicide, there are approximately 25 suicide attempts.⁸ For youth aged 15 to 24, as many as 200 attempts may occur for every death.⁹ Suicidal thoughts are more common. In 2017, for example, an estimated 9.8 million adults nationally reported experiencing suicidal thoughts, but far fewer – 2.8 million adults – made suicide plans, while 1.3 million adults attempted suicide.¹⁰ Beyond the profound impact on the person, family, community, and society, suicide poses an estimated economic cost of \$93.5 billion in lost productivity and medical expenses in the U.S.¹¹ In California, suicide resulted in an average of \$1,085,227 per death in lost productivity and medical expenses in 2010.¹² This does not include the cost of other suicidal behavior, such as suicide attempts that did not result in death.

Suicide has emerged as a public health emergency in need of innovation across multiple levels of prevention because of historically intractable rates.¹³ A public health approach is suitable to meet the challenge.¹⁴ This approach seeks to increase the health of the community in order to reduce the risk experienced by each person and, likewise, to increase the health of each person to reduce risk in the community.¹⁵ Health, therefore, is determined by the physical, psychological, cultural, and social environments in which people live, work, and go to school.¹⁶ The Public Health Model guides this approach, and it involves the following: defining the problem; identifying the factors that increase or lower risk; developing and evaluating prevention interventions; and implementing interventions and disseminating results to increase the use of effective interventions.¹⁷ See the figure on the next page. The Public Health Model is a key feature of the statewide strategic suicide prevention plan detailed in this document.



Public Health Model adapted from WHO Preventing Suicide: A Global Imperative

California's Strategic Plan for Suicide Prevention

The first half of California's Strategic Plan for Suicide Prevention outlines the strategic aims, goals, and actions needed to prioritize suicide prevention efforts across the state over the next five years, with the ultimate goal of no loss of life to suicide. These pages detail the tactics, or "how to" steps, that can help California communities effectively prevent suicide given the latest in understanding of best practices. The second half of the plan details shared strategic understanding of terms, theory, challenges, and evidence to support the coordinated delivery of suicide prevention efforts.

Strategic Aims, Goals, and Action

The following plan establishes a foundation of suicide prevention directed by best practices for the benefit of state and local partners. The goal is ambitious, but achievable – to ensure no life is lost to suicide in California. Suicide prevention is a responsibility shared among private and public partners, and likewise, efforts must be driven by private and public data and resources, including human and fiscal assets. State funding should support key areas outlined in the action steps that follow, which include establishing state leadership, delivering technical assistance, developing guidance, and fortifying and expanding data collection and reporting systems. However, to ensure the sustainability efforts, other public and private assets must be leveraged and continuously pursued.

Four strategic aims and 12 goals serve as a roadmap to align local efforts with state priorities in delivering best practices in suicide prevention. Local communities can start now to identify local health and behavioral health leaders, build coalitions, and begin to identify data and information to describe the problem of suicidal behavior in their communities. Once the problem is described, communities can continue the public health model by identifying risk and protective factors, developing interventions and conducting evaluation, and disseminating effective practices.

Plan Components

This plan serves as strategic guidance to equip local communities with information on best practices and areas of focus with the greatest potential for preventing suicide in their communities. The plan is organized using the following components:

- **Strategic aims** are broad, long-term goals to reduce suicidal behavior.
- **Goals** accompany each strategic aim to serve as a roadmap for governments, community organizations, providers, and other partners to focus suicide prevention efforts using best-practice approaches or interventions. These are discussed in greater detail in the Best Practice in Suicide Prevention section of this plan.
- **Key action partners** are identified to support state and local leaders to advance each goal.
- **Strategies** for achieving the goal at both the state and local levels are included under each goal.
- **Short-term targets and long-term outcomes** are identified under each goal. Measuring incremental steps and progress toward reaching each goal, while monitoring suicide data, will be critical.¹⁸
- **Action to implement state strategies** are next steps the state should take to support local implementation and statewide advancement of strategies listed after each goal.

Plan Quick View

California's Strategic Plan for Suicide Prevention is framed by four strategic aims and 12 goals. Each goal statement embeds suicide prevention strategies and approaches with the greatest potential to prevent suicide in communities across the state. See the Best Practices in Suicide Prevention section of this plan for a detailed description of the evidence to support the effectiveness of each strategy.

Strategic Aim 1:

Establish infrastructure to support suicide prevention strategies

- Goal 1: Enhance visible leadership and networked partnerships
- Goal 2: Strengthen management of suicide prevention resources
- Goal 3: Advance data monitoring and evaluation

Strategic Aim 2:

Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

- Goal 4: Create safe environments by reducing access to lethal means
- Goal 5: Increase resiliency and help-seeking
- Goal 6: Increase connectedness between people, family members, and community
- Goal 7: Increase safe reporting of suicide and healthy social media use

Strategic Aim 3:

Increase early identification of suicide risk and access to services based on risk

- Goal 8: Increase detection and screening to connect people to services
- Goal 9: Deliver a continuum of crisis services within and across counties

Strategic Aim 4:

Advance suicide-specific services and supports

- Goal 10: Deliver best practices in care targeting suicide risk
- Goal 11: Ensure continuity of care and follow-up after suicide-related services
- Goal 12: Expand support services following a suicide loss

Strategic Aim 1: Establish infrastructure to support suicide prevention strategies

Goal 1: Enhance visible leadership and networked partnerships

Strategic Aim 1: Establish infrastructure to support suicide prevention strategies

Short-term Target

- By 2025, state leadership is advancing suicide prevention as a public health priority, and all counties have leaders and coalitions engaged in suicide prevention efforts

Long-term Outcome

- Increase awareness and sustainability of suicide as a preventable public health priority

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders and providers• Indigenous and traditional healers• Youth leaders• Parents and caregivers• School, college, and university staff	<ul style="list-style-type: none">• Business and nonprofit leaders• Community service providers, especially providers serving vulnerable populations• Faith-based leaders• Criminal and juvenile justice leaders• Suicide prevention subject matter experts

State Strategies to Support Goal

- Establish centralized, visible state-level leadership to provide strategic guidance, deliver technical assistance, develop and coordinate trainings, monitor data, and conduct state-level evaluation, and disseminate information to advance statewide progress.
- Engage private and public partners to advance suicide prevention efforts with strategic planning and dissemination of best practices in their respective sectors.

Local Strategies to Support Goal

- Establish leadership to provide clear direction for suicide prevention efforts and prioritize goals with maximal impact. Suicide prevention leadership may come from a coalition, task force, or health or behavioral health agencies or organizations.

- Identify leaders who can champion suicide prevention as a public health priority. Leaders drive progress, develop and sustain relationships with partners, and help focus attention on suicide prevention as a core mission when faced with competing priorities.
- Hold regularly scheduled meetings to convene stakeholders, prioritize suicide prevention activities based on data and community input, leverage resources to build capacity across systems and communities/regionally, and expand services based on effectiveness.
- Formalize a coalition of private and public partners to advance suicide prevention efforts by being an “action arm” to local leaders.¹⁹ Private and public leaders should be brought together to leverage their influence to champion efforts prioritized in their own sectors.²⁰ Create sector-specific or strategy-specific subgroups to focus expertise within suicide prevention coalitions, and keep members energized and engaged.²¹ Provide consistent and predictable infrastructure by local leadership to the coalition, which may include logistical support, strategic guidance, and technical assistance.²²

Goal 2: Strengthen management of suicide prevention resources

Strategic Aim 1: Establish infrastructure to support suicide prevention strategies

Short-term Target

- By 2025, all counties are working to prioritize suicide prevention as a public health challenge, and are implementing suicide prevention initiatives, including establishing a dedicated website listing local suicide prevention resources, forming coalitions, and creating suicide prevention strategic plans.

Long-term Outcome

- Increase in private and public assets to expand effective suicide prevention activities

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • Health, public health, and behavioral health leaders and providers • Indigenous and traditional healers • Parents and caregivers • School, college, and university staff • Youth leaders 	<ul style="list-style-type: none"> • Business and nonprofit leaders • Community service providers, especially providers serving vulnerable populations • Faith-based leaders • Criminal and juvenile justice leaders • Suicide prevention subject matter experts

State Strategies to Support Goal

- Advance the development and management of suicide prevention resources in communities across California, and support capacity building to use best practices in suicide prevention by disseminating guidance and resources.
- Identify and implement opportunities to integrate suicide prevention strategies across systems and programs.

Local Strategies to Support Goal

- Develop a local suicide prevention plan and implementation strategy to prevent suicidal behavior across the lifespan and to address the goals outlined in the state's strategy, in addition to addressing local needs. Funding allocated to local behavioral health departments under the Mental Health Services Act can be used for the purposes of suicide prevention planning, as well as developing and implementing strategies.
- Map local assets across sectors to coordinate resources and align funding priorities, especially if partners have data that can demonstrate how investments in specific suicide prevention strategies could lead to improved outcomes and cost savings in other areas. Assets may be programs or features of the community, such as safe and welcoming community spaces, parks, or centers. Assets can be mobilized through planning processes that identify often underutilized community strengths, such as Asset-Based Community Development strategies.²³
- Document the roles and responsibilities of each partner, and any data or funding streams associated with each partner and their affiliation. Each partner has a role to play, and all partners bring potential for innovating common practices.
- Identify opportunities to integrate suicide prevention strategies into services already being delivered through local settings, systems, and programs. For example, train community health workers to recognize warning signs of suicide and equip them with resources to encourage people to follow safe storage practices - such as how to dispose of unused medications - and directly connect people at risk to care or crisis services.
- Leverage partnerships through a coalition (see Goal 1) to identify shared prevention goals across diverse settings and communities, such as education, child welfare, social services, health care, and justice settings. These partners may have shared goals with suicide prevention for reducing risk and increasing protective factors, such as creating safe and active communities to reduce social isolation. All can be leveraged to reduce suicidal behavior and meet other goals for health and wellness promotion.

Goal 3: Advance data monitoring and evaluation

Strategic Aim 1: Establish infrastructure to support suicide prevention strategies

Short-term Target

- By 2025, 100 percent of all suicide deaths are electronically entered into the California Violent Death Reporting System and communities are using timely aggregated data to strengthen suicide prevention strategies

Long-term Outcome

- Increase in the use of data to guide suicide prevention state and local policy and planning, resource management, and investment

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders and providers• Indigenous and traditional healers• Youth leaders• Parents and caregivers• School, college, and university staff	<ul style="list-style-type: none">• Business and nonprofit leaders• Community service providers, especially providers serving vulnerable populations• Faith-based leaders• Criminal and juvenile justice leaders• Suicide prevention subject matter experts

State Strategies to Support Goal

- Establish centralized, electronic reporting systems to capture data related to suicide deaths and suicidal behavior, with uniform coding procedures.
- Develop a data monitoring and evaluation agenda on suicide deaths and suicidal behavior, including data elements documenting interrupted or aborted suicide attempts and crisis service interventions (“save data”) that resulted in the de-escalation of suicidal desire or intent.
- Standardize policies and procedures for investigating and reporting suicide as a cause of death, including uniform definitions of suicide, as well as protocols for working with suicide loss survivors and informing health officials in the context of a suicide cluster. Include clear requirements for how cause of death is determined, how investigations are conducted, and how information is reported, and by whom, within a certain time following death. Include training on methods for minimizing misclassification and accelerating timely reporting.

Local Strategies to Support Goal

- Use local data and information to define the problem of suicidal behavior, identify factors that increase or lessen risk for suicide, develop interventions, conduct evaluations, and disseminate effective practices that prevent suicide.
- Use suicide death and attempt data to evaluate the proportion of suicidal behavior that results in death. Use results to identify high-risk groups, target them with selective prevention strategies, and focus resources on specific lethal means restriction strategies.
- Use death review team models for clinical and forensic review of suicide deaths. Team members should include representatives of coroners and medical examiners, law enforcement, subject matter experts, and others with legal access to confidential information. Use data compiled by the team to support prevention goals using the Public Health Model.
- Partner with coroners and medical examiners to identify barriers and deliver support for electronic reporting of suicide death data into California's Violent Death Reporting System. Develop a method for accessing data for improving suicide prevention strategies, including establishing policies and procedures for protecting privacy.
- Use community surveys to fill data gaps. For example, people with non-fatal, self-directed violence may not seek medical attention following the injury, thereby reducing the number of such reports.²⁴ Community surveys could be used to supplement data.
- Build relationships with local colleges and universities and identify capacity for research.

Action to Implement State Strategies

Goal 1:	Enhance visible leadership and networked partnerships
State Strategy	Action to Implement Strategy
Establish centralized, visible state-level leadership to provide strategic guidance, deliver technical assistance, develop and coordinate trainings, monitor data, and conduct state-level evaluation, and disseminate information to advance statewide progress.	<p>By July 1, 2021, the State should create the Office of Suicide Prevention and position the office under the California Health and Human Services Agency.</p> <p>By December 31, 2021, the Office of Suicide Prevention should develop a plan to facilitate regional quarterly meetings across the state to share resources, best practices, and lessons learned in developing strategies to deliver a continuum of crisis services to prevent suicidal behavior.</p> <p>By July 1, 2022, the Office of Suicide Prevention should form a task force of subject matter experts for Strategic Aims 2, 3, and 4 to create a research and policy agenda as described under each goal.</p> <p>By July 1, 2022, the Office of Suicide Prevention should develop a strategy for leveraging federal grant and block grant funding and private investment in suicide prevention strategies.</p> <p>By July 1, 2023, the Office of Suicide Prevention should host and maintain an online clearinghouse to support implementation of best practices and technical assistance.</p>
Engage private and public partners to advance suicide prevention efforts with strategic planning and dissemination of best practices in their respective sectors.	<p>By July 1, 2021, the State should create the California Suicide Prevention Council and appoint councilmembers.</p> <p>By December 31, 2021, the California Suicide Prevention Council should hold its first meeting and develop a strategic work plan. The work plan should include how the council will support the state strategies outlined in this plan.</p> <p>By July 1, 2022, the California Suicide Prevention Council should form sector-specific or strategy-specific subgroups to focus expertise within the council and develop guidance to support suicide prevention efforts in specific sectors.</p>

Goal 2:	Strengthen management of suicide prevention resources
State Strategy	Action to Implement Strategy
<p>Advance the development and management of suicide prevention resources in communities across California, and support capacity building to use best practices in suicide prevention by disseminating guidance and resources.</p>	<p>By July 1, 2021, the State should create incentives for local suicide prevention planning and implementation, including offering grants to support capacity building to deliver best practices prioritized in the state's plan.</p> <p>By July 1, 2021, the State should amend existing legislation requiring public schools with students in grades seven through 12 to develop a suicide prevention policy by including a provision of oversight by the Department of Education. The amendment should require schools to submit policies to the department for review and dissemination, and the department should deliver technical assistance and support to schools without policies. The department also should examine barriers to suicide prevention identified by schools – including liability issues, privacy laws, security measures, and legal requirements for parental consent – and develop recommendations to address them.</p> <p>The Department of Education should evaluate the effectiveness of current school policies and revise its model policy based on best practice. In addition, the department should develop a strategy for evaluating policies on an ongoing basis, through metrics such as reductions in suicidal behavior, increases in connection to services, and increases in students and school personnel seeking help.</p> <p>By July 1, 2021, the State should expand the requirement that public schools develop suicide prevention policies to include colleges and universities.</p> <p>By July 1, 2022, the Office of Suicide Prevention should develop and deliver a process for disseminating information to support local suicide prevention planning and implementation, which may include regional learning collaboratives to share resources and data, best practices, and lessons learned in delivering local suicide prevention strategies.</p>
<p>Identify and implement opportunities to integrate suicide prevention strategies across systems and programs.</p>	<p>By July 1, 2022, the Office of Suicide Prevention and the California Suicide Prevention Council should develop and implement a process for collaborating with public and private partners to integrate suicide prevention strategies across statewide programs and initiatives.</p>

Goal 3:	Advance data monitoring and evaluation
State Strategy	Action to Implement Strategy
<p>Establish centralized, electronic reporting systems to capture data related to suicide deaths and suicidal behavior, with uniform coding procedures.</p>	<p>By July 1, 2021, the State should authorize counties to utilize death review team models to evaluate local suicidal behavior trends, circumstances, and outcomes to inform and strengthen local prevention strategies.</p> <p>By July 1, 2021, the State should create incentives for schools to regularly participate in the California Healthy Kids Survey to monitor trends in suicidal behavior among students, including allocating additional resources to create reports on student suicidal behavior that are specific to each school.</p> <p>By December 31, 2021, the State, with leadership from the Department of Public Health, should expand the existing California Violent Death Reporting System (CalVDRS) to more counties to collect and analyze local and state suicide data by delivering technical assistance to local coroners and medical examiners. The assistance should enhance the timely and electronic reporting of suicide deaths and their circumstances – including contributing factors and the specific location of death if outside the home – to help identify and fortify the safety of sites used by people to die by suicide.</p> <p>The State should invest additional resources to fund technical assistance to increase the participation by coroners, medical examiners and law enforcement agencies in the CalVDRS to provide more detailed information on circumstances surrounding violent deaths, including suicide.</p> <p>By January 1, 2022, the State, with leadership from the Department of Public Health and the Department of Health Care Services, should identify additional data elements to be collected via the California Health Interview Survey to include data on suicide risk and protective factors to monitor suicidal behavior across the state.</p> <p>By July 1, 2023, the State, including private and public partners, should develop and implement a strategy to improve the standardization of coding and reporting of suicidal behavior,</p>

Goal 3:	Advance data monitoring and evaluation
State Strategy	Action to Implement Strategy
	<p>including the development of guidelines for determining suicidal intent. The state also should develop a plan to deliver training and technical assistance to hospital representatives to improve the identification, coding, and reporting of suicidal behavior for people seen in emergency departments and admitted to hospitals.</p> <p>By December 31, 2023, the State, including private and public partners, should create a mechanism for centralized and electronic reporting the number of people who are screened for suicide risk in hospitals and emergency departments, and data documenting how people who were positively identified at various levels of risk were triaged into services. For example, data entered into electronic health records could be extracted and aggregated prior to its submission to a centralized database.</p>
<p>Develop a data monitoring and evaluation agenda on suicide deaths and suicidal behavior, including data elements documenting interrupted or aborted suicide attempts and crisis service interventions (“save data”) that resulted in the de-escalation of suicidal desire or intent.</p>	<p>By December 31, 2021, the Office of Suicide Prevention should create a task force to develop a data monitoring and evaluation agenda on suicidal behavior, including data elements documenting interrupted or aborted suicide attempts and crisis service interventions that resulted in the de-escalation of suicidal desire or intent. The task force should identify opportunities for expanding research exploring community-defined practices that reduce suicide risk in diverse cultural groups and disseminate findings directly to communities affected and the public. The task force should include in the agenda guidance for local program evaluation and should identify measures to monitor state-level outcomes.</p> <p>By July 1, 2023, the task force should develop for the Governor and Legislature a proposal to create a centralized, electronic database and reporting standards to capture data on interrupted or aborted suicide attempts and crisis service interventions that resulted in the de-escalation of suicidal desire or intent. The data must include the intervention used and should include the type of services referred and the duration between incident and entry into services. Data sources include, but are not limited to, first responders, emergency and health care practitioners and providers, crisis service providers, and bridge and transportation</p>

Goal 3:	Advance data monitoring and evaluation
State Strategy	Action to Implement Strategy
	<p>representatives. The proposal must include an estimate for costs associated with the centralized database.</p>
<p>Standardize policies and procedures for investigating and reporting suicide as a cause of death, including uniform definitions of suicide, as well as protocols for working with suicide loss survivors and informing health officials in the context of a suicide cluster. Include clear requirements for how cause of death is determined, how investigations are conducted, and how information is reported, and by whom, within a certain time following death. Include training on methods for minimizing misclassification and accelerating timely reporting.</p>	<p>By July 1, 2023, the Office of Suicide Prevention should form a task force to develop and disseminate best practices in suicide death investigation procedures, including guidance for coroners and medical examiners for documenting behavioral issues, hospitalizations, medications, histories of suicidal behavior, and family behavioral health history.</p> <p>Guidance should include methods for sharing data with local or state death review teams with the goal of identifying opportunities for improvement in prevention strategies. Guidance should include guidelines for coroners and medical examiners for identifying and reporting sexual and gender diversity of people who die by suicide and include recommendations for any necessary modifications to existing reporting systems to enable data collection on suicides by sexually and gender diverse people.</p>

Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

Goal 4: Create safe environments by reducing access to lethal means

Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

Short-term Target

- By 2025, 50 percent increase in the distribution of lethal means safety products, education, and information

Long-term Outcome

- Reduce suicidal behavior using highly lethal means, such as firearms

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders and providers• Tribal community members• In-home service providers• Youth leaders• Parents and caregivers• Law enforcement leaders	<ul style="list-style-type: none">• Suicide prevention organizations• Gun and gun shop and range owners• Gun and shooting clubs• Pharmacy administrators• Community-based organizations and nonprofits• Aging service providers• Transportation leaders

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of creating safe environments by reducing access to lethal means.
- Implement a method for monitoring state-level trends in lethal means for in suicidal behavior and a plan for safely disseminating trend information and emerging best practices for reducing means.
- Deliver technical assistance to expand methods for reducing access to lethal means for the purposes of preventing suicidal behavior.

- Disseminate information regarding federal funding available to support suicide barriers in the new design or re-design of bridge and other sites to prevent deaths at sites where suicides occur.

Local Strategies to Support Goal

- Use the Public Health Model to evaluate risk and identify the methods of suicidal behavior used by community members and by specific demographic and cultural groups to guide the development of focused prevention efforts. Once identified, develop tailored means restriction strategies and evaluate impact.
- Promote safe medication disposal methods in the community or through pharmacies and other health care practitioners and providers, including activities such as “take back” campaigns led by local public health departments that help people dispose of unused or expired medications. Partner with local pharmacies to increase the availability of methods to dispose of unused medication and highlight suicide and overdose prevention resources for people filling prescriptions.
- Disseminate information to local gun shop and range owners to increase awareness of suicide prevention efforts, suicide warning signs, and available resources. Partner with local firearm safety trainers to incorporate suicide prevention awareness into trainings. Invite local gun shop and range owners to join local coalitions. Partner with law enforcement to guide dissemination of lawful options for temporarily transferring firearms for storage in times of suicide crisis or use of gun restraining orders.
- Disseminate information through the local health department to community partners about medication assisted therapy options, such as naloxone, in the event of an opioid overdose.
- Form a workgroup of community members, first responders, transportation representatives, coroners and medical examiners, and crisis service providers to identify specific sites in the community used frequently for suicide, or those that provide the opportunity for suicide. These sites can be in the built environment or natural sites. Common types of sites include buildings, bridges, and train railways. Characteristics communities should consider in identifying sites are places that provide the opportunity for a person at risk to fall from a height and sites from which falling would place a person in front of a moving vehicle, such as a train. More than one suicide at a particular site should raise safety concerns.

- Once sites are identified, develop and implement plans to construct barriers to deter or prevent falling and install signs that list crisis services resources, such as suicide prevention hotline information, and provide positive, life-affirming messages.
- Create memorandums of understanding or other agreements with local bridge and rail authorities, first responders, and crisis services providers to develop a method of collecting data documenting events in which people were prevented from falling, the type of resources to which people were referred, and the outcome of connection to those resources. Include reporting requirements, such as biannual or quarterly reports.

Goal 5: Increase resiliency and help-seeking

Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

Short-term Target

- By 2025, 10 percent increase in behavioral health services utilization

Long-term Outcome

- Reduce the number of people unserved and underserved for behavioral health needs

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • Health, public health, and behavioral health leaders and providers • Parents and educators • Suicide prevention organizations • Faith-based leaders 	<ul style="list-style-type: none"> • Community-based organizations and nonprofits • Youth leaders • Business leaders • Tribal leaders • Aging service providers

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of increasing resiliency and help-seeking.
- Integrate social-emotional learning programs into private and public education curricula for the purposes of strengthening communication and problem-solving skills, emotional regulation, and conflict resolution skills among children and youth.

Local Strategies to Support Goal

- Identify community needs for managing stressors and building resiliency, which may include coping skills, critical thinking, stress management, conflict resolution, and problem-solving skills.
- Identify barriers community members face in seeking services for behavioral health needs, and develop strategies for making services more accessible, convenient, and culturally respectful to increase the likelihood people will pursue and stay connected to such services.
- Expand outreach and engagement strategies to promote behavioral health and community services and resources.
- Integrate activities into community-based services that increase life skills, including mindfulness practices, critical thinking, stress management, conflict resolution, problem-solving, and coping skills; tailor activities based on age group and setting, and according to how different groups experience and mitigate stress. Cultural models of suicide can be helpful in understanding how culture affects the experiences of stressors, the cultural meaning of stressors, and how different cultures express suicidal behavior.²⁵
- Expand services to increase mental health literacy, encourage people to seek help for health and behavioral health needs, and promote messages of hope that lives can be saved from suicide.
- Develop a network of peer support providers to help people navigate health care and behavioral health care systems. “Peer” can include a person with lived experience, a survivor of suicide loss or attempt, or a person from a peer group, such as veterans, first responders, or students. Build peer capacity among students to communicate distress and have clear and easy pathways to caring adults who can help them navigate their options.
- Partner with community organizations and business to expand awareness of suicide warning signs and prevention resources. Coordinate suicide prevention awareness campaigns with other social marketing campaigns designed to reduce mental health stigma and discrimination and reduce relevant public safety threats, such as misuse of medication or unsafe gun storage practices.
- Use suicide loss and attempt survivors to develop and deliver strategies that can increase the number of people seeking help by reducing stereotypes about people who sought out services and benefitted from those services. For example, people with lived

experience can be effective service navigators, helping others understand the service delivery system and reducing confusion or misconceptions. Methods are especially meaningful when they include cultural congruency between people with lived experience and a target audience, such as youth helping youth or veterans helping veterans.

- Create memorandums of understanding defining responsibilities and uniform suicide prevention performance metrics between organizations and support services.

Goal 6: Increase connectedness between people, family members, and community

Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

Short-term Target

- By 2025, 20 percent increase in community-based services intended to reduce social isolation and strengthen relationships between people and their families and caregivers

Long-term Outcome

- Reduce exposure and experience of traumatic adverse events

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • Health, public health, and behavioral health leaders and providers • Parents and educators • Suicide prevention organizations • Youth leaders 	<ul style="list-style-type: none"> • Community-based organizations • Business leaders • Tribal leaders • Aging service providers • Faith-based leaders

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of increasing connectedness between people, family members, and community.
- Align efforts and investments to address multiple forms of violence that may share risk and protective factors with suicide, including strategies for reducing trauma in early childhood.

Local Strategies to Support Goal

- Increase services focused on building positive attachments between children, youth, and adults, their families, and social networks in their community to increase sense of belonging, strengthen sense of identity and personal worth, and provide access to larger sources of support. Social networks can be found in schools, faith-based communities, cultural centers, and other community-based organizations.
- Tailor strategies to be responsive to needs based on age and culture. For example, create peer support groups, led by veterans or active duty members of the military, which allow veterans to safely share their experiences; and disseminate talk-based warmline phone numbers targeting older adults to reduce feelings of isolation and loneliness, and use communication methods relevant to an older population, such as advertising in health care settings or through traditional media.
- Promote a culture free of stigma and discrimination by allowing for an open dialogue about mental health and mental health resources, and by delivering supportive messages of hope and recovery for people with behavioral health needs. Establish policies and methods for enforcement to create cultures that support healthy lifestyles and environments that are affirmative and that prevent violence, including bullying and discrimination.
- Identify opportunities to integrate suicide prevention strategies into services intended to reduce other forms of violence, such as child and elder maltreatment. These forms of violence may share risk and protective factors with suicidal behavior. For example, reducing interpersonal stress and teaching conflict resolutions skills among at-risk families has the potential to increase a sense of connectedness and protects against suicide.
- Partner with community-based organizations to build and promote opportunities for volunteerism to increase connectedness and a sense of purpose.

Goal 7: Increase safe reporting of suicide and healthy social media use

Strategic Aim 2: Minimize risk for suicidal behavior by promoting safe environments, resiliency, and connectedness

Short-term Target

- By 2025, all counties are delivering training for local media partners on safe reporting of suicides

Long-term Outcome

- Increase the number of California-based media outlets using safe reporting practices

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders and providers• Youth leaders• Parents and educators• Suicide prevention organizations• Behavioral health media consultants	<ul style="list-style-type: none">• Journalists and news organizations• Representatives of the technology industry• Representatives from schools of journalism and high school journalism programs• Local spokespeople• Entertainment industry leaders

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of increasing safe reporting of suicide and social media use.
- Increase awareness of safe reporting on suicides and training for journalism and media partners on how to safely write about suicide.
- Develop curricula for high school and college journalism students to teach best practices for writing about suicide through various forms of media and entertainment.
- Identify and disseminate best practices for using and consuming social media to improve wellbeing, destigmatize mental illness, and increase help-seeking for behavioral health services.

Local Strategies to Support Goal

- Identify media and entertainment industry partners and deliver training on best practice guidelines for reporting about suicide. Disseminate information found online at <http://reportingonsuicide.org/> and <http://suicidepreventionmessaging.org/> to members of the media – reporters, editors, and producers – regarding how risk is conferred and to improve understanding of guidelines supporting suicide prevention on a broad scale.
- Partner with members of media to disseminate information about resources, encourage people to seek help for behavioral health needs, and reduce stigma and discrimination that may prevent people from accessing services and supports. Entertainment media include film, television, music, and theater.
- Deliver training to parents to support healthy use of social media by youth. Disseminate the American Association of Suicidology tip sheet for parents and providers, available at <https://www.suicidology.org/LinkClick.aspx?fileticket=9BI5g4TXz54%3D&portalid=14>.
- Disseminate information about how suicide risk can be expressed by people on various social media sites and highlight social media resources for identifying and reporting concerns about content. Most social media sites now have a method for reporting content that raises alarms.
- Integrate into public campaigns and health and mental health curriculum in schools best practices for developing healthy social media habits and using social media in a way that promotes connectedness to reduce isolation.
- Create communication strategies for use in the event of a suicide loss – including pre-existing agreements with media partners – to minimize the circulation of misinformation. Include a formal strategy for managing information on the most used social media sites and monitor social media posts by others related to the suicide loss.

Action to Implement State Strategies

Goal 4:	Create safe environments by reducing access to lethal means
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of creating safe environments by reducing access to lethal means.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 2.
Implement a method for monitoring state-level trends in lethal means for in suicidal behavior and a plan for safely disseminating trend information and emerging best practices for reducing means.	By December 31, 2021, the Office of Suicide Prevention should enter into data use agreements to receive suicide-related data from state departments to monitor the use of lethal means in suicidal behavior and evaluate trends. Based on information, the office should use the data to tailor technical assistance resources.
Deliver technical assistance to expand methods for reducing access to lethal means for the purposes of preventing suicidal behavior.	By July 1, 2022, State, with leadership from the Department of Public Health, should develop and implement a technical assistance strategy to expand information and availability of methods that can prevent injury due to suicidal behavior and death by suicide, including gun locks, gun and medication safes, devices to dispose of unused medication, and medications to counteract overdose, such as naloxone for opioid overdose.
Disseminate information regarding federal funding available to support suicide barriers in the new design or re-design of bridge and other sites to prevent deaths at sites where suicides occur.	<p>By December 31, 2022, the Office of Suicide Prevention should create an online clearinghouse of strategies and resources for reducing access to lethal means, including information on available private and public funding. The online clearinghouse should include methods to accelerate dissemination and implementation of best practices, such as quick factsheets and “how to” guides. The online clearinghouse should include information on new approaches to reducing access to lethal mean as they emerge.</p> <p>By December 31, 2023, the Office of Suicide Prevention should form a task force to review and make recommendations for modifying buildings, bridges, and other structures if such modifications are to prevent suicide at identified locations. The office should partner with the California Coastal Commission, the Office of Historic Preservation, transportation leaders, and others to address “line of sight” and other aesthetic concerns that may impede modifications that improve safety.</p>

Goal 5:	Increase resiliency and help-seeking
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of increasing resiliency and seeking services for behavioral health needs.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 2.
Integrate social-emotional learning programs into private and public education curricula for the purposes of strengthening communication and problem-solving skills, emotional regulation, and conflict resolution skills among children and youth.	By July 1, 2024, the State, with leadership from the Department of Education, the California State Board of Education, and Instructional Quality Commission, should develop standards for social emotional learning and require implementation of such standards in schools.

Goal 6:	Increase connectedness between people, family members, and community
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of increasing connectedness between people, family members, and community.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 2.
Align efforts and investments to address multiple forms of violence that may share risk and protective factors with suicide, including strategies for reducing trauma in early childhood.	By July 1, 2022, the State, with leadership from the Department of Public Health and private and public partners, should conduct an environmental scan of population-based universal violence prevention strategies and programs across the state, including suicide prevention programs, as well as those that address shared risk and protective factors for multiple forms of violence.

Goal 6:	Increase connectedness between people, family members, and community
State Strategy	Action to Implement Strategy
	<p>By December 31, 2022, the State, with leadership from the Department of Public Health and private and public partners, should develop recommendations to help communities increase community cohesion and safety, especially for vulnerable groups, and highlight areas of California where programs are making an impact. The effort should focus on ways to increase key protective factors, including connectedness, positive social norms, resiliency, and economic opportunity.</p> <p>By July 1, 2023, the State, with leadership from the Department of Public Health and private and public partners, should identify a common set of measures and indicators that could be used by programs addressing violence prevention to enhance alignment, track progress, and improve understanding of needs and gaps statewide.</p>

Goal 7:	Increase safe reporting of suicide and healthy social media use
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of Increasing safe reporting of suicide and social media use.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 2.
Increase awareness of safe reporting on suicides and training for journalism and media partners on how to safely write about suicide.	By July 1, 2022, the Office of Suicide Prevention should create a task force with media and journalism outlets and organizations that publish journalism ethics codes to develop a process for ensuring the use of safe reporting on suicides. This effort should produce state standards for safe reporting and messaging about suicide in the media, guidance for partnering with media and entertainment industry representatives, and a strategy for dissemination, including incentives for media outlets to adhere to standards.

Goal 7:	Increase safe reporting of suicide and healthy social media use
State Strategy	Action to Implement Strategy
	By December 31, 2023, the task force should be building relationships with media industry partners and developing and disseminating resources to promote depictions of positive storylines of mental health and suicide prevention. Resources should include guidance to help media writers and producers accurately and safely portray suicidal behavior. Potential actions the task force could take for industry partners include fact-checking and research, background information about suicide risk and protective factors and warning signs, and advice on tone and use of language in scripts.
Develop curricula for high school and college journalism students to teach best practices for writing about suicide through various forms of media and entertainment.	By July 1, 2024, the Office of Suicide Prevention should form a task force to create a curriculum for reporting and writing about suicide deaths in the media to be delivered in high school, college, and graduate journalism programs.
Identify and disseminate best practices for using and consuming social media to improve wellbeing, destigmatize mental illness, and increase help-seeking for behavioral health services.	By July 1, 2024, the State, including private and public partners, should develop a process for disseminating information and resources on the appropriate and safe use of social media, tailored to age-group and setting, and information and resources on how parents and caregivers can support the healthy use of social media.

Strategic Aim 3: Enhance early identification of suicide risk and increase access to services based on risk

Goal 8: Increase detection and screening to connect people to services based on suicide risk

Strategic Aim 3: Enhance early identification of suicide risk and increase access to services based on risk

Short-Term Target

- By 2025, all people served in health care settings are routinely screened for suicide using uniform best practices in suicide risk assessment and management

Long-Term Outcome

- Decrease in suicidal behavior and increase in connection to services

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders• Community health workers• Indigenous and traditional healers• Youth leaders	<ul style="list-style-type: none">• Workplace supervisors and leaders• Community service providers, especially providers serving vulnerable populations• Faith-based leaders• School, college, and university staff• Parents and caregivers

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of increasing detection and screening to connect people to services based on suicide risk.
- Expand resources to support health care practitioners and providers increase access and linkage to behavioral health services for people identified as needing such services.
- Adopt the Zero Suicide Initiative within health care systems.
- Increase standardized training in best practices in suicide risk assessment and management to health care practitioners and providers and enhance uniform suicide risk assessment and management in health care settings, in alignment with Joint Commission guidelines and the Zero Suicide Initiative.

- Identify opportunities for technology in health care systems to advance suicide risk assessment and management application, use, and investment.

Local Strategies to Support Goal

- Deliver suicide prevention training to people in positions to identify warning signs of suicide. Support youth gatekeepers by identifying trusted adults to help them with next steps once a student is identified as at risk. Provide people the opportunity to reinforce knowledge and skills acquired during training through periodic booster sessions. Build capacity for suicide prevention training across systems using train-the-trainer models.
 - Consider the intensity of training needed and offer a variety of trainings to expand capacity to meet differences in demand. For example, in a school setting, teachers, administrators, and other school personnel might receive brief trainings on suicide prevention awareness. Selected teachers, especially those who lead youth groups, and counselors might receive intensive trainings on delivering brief interventions.
- Screen people seen in health and behavioral health care settings for suicide risk and deliver best practices in suicide risk assessment and management for people who positively screen for risk.
 - The Joint Commission recommended screening and assessment tools include the following: Ask Suicide Screening Toolkit (ASQ) by National Institute of Mental Health; the Columbia Suicide Severity Rating Scale (C-SSRS) Triage Version; Patient Health Questionnaire 9 (PHQ-9) Depression Scale; Suicide Behavioral Questionnaire Revised; Scale for Suicidal Ideation-Worst; and the Beck Scale for Suicide Ideation.²⁶
 - Suicide screenings can follow positive results on other screening tools. For example, screening specific to suicide risk should follow positive screens for depression, anxiety, trauma, physical pain, and problem alcohol, drug use, and eating. Comprehensive suicide risk assessments follow screening.
- Integrate best practices in suicide risk assessment and management in health care settings and workflows and clarify methods for billing for services. Create uniform policies and procedures to routinize screening, assessments, and decision-making.
- Deliver training to key action partners for conducting suicide screening in community-based settings when a person is identified as exhibiting warnings signs or

communicating a desire to die. The Columbia-Suicide Severity Rating Scale has been adapted to meet the needs of diverse settings and populations, and can be accessed for free here: <http://cssrs.columbia.edu/>.

- Deliver training for first responders and other personnel patrolling or monitoring community sites used for suicidal behavior, such as bridges and railways. The training should include how to identify warning signs, use de-escalation techniques, and disseminate information on local suicide prevention resources, including crisis hotline numbers.

Goal 9: Deliver a continuum of crisis services within and across counties

Strategic Aim 3: Enhance early identification of suicide risk and increase access to services based on risk

Short-Term Target

- By 2025, 20 percent increase in utilization of crisis services to reduce suicide risk and connect people to community-based services

Long-Term Outcome

- Decrease in suicidal behavior among people accessing crisis services

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • First responders • Health, public health, and behavioral health leaders and providers • Crisis services providers and administrators • Suicide prevention organizations 	<ul style="list-style-type: none"> • Indigenous and traditional healers • Parents and caregivers • School, college, and university staff • Workplace supervisors and leaders • Youth leaders

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of delivering a continuum of crisis services within and across counties.
- Develop and implement a strategy to coordinate the delivery of crisis services, including an assessment of current crisis services infrastructure and private and public funding for services.

- Develop a state accreditation program for suicide and crisis hotlines operated in the state, including standards for training and core competencies for call responders, protocols for performance and quality assurance monitoring, and procedures for making referrals to services, including emergency services.

Local Strategies to Support Goal

- Evaluate the continuum of crisis services available through private and public resources and identify continuum gaps and potential funding sources.
- Promote the use of crisis services as alternatives to hospitalization and as a resource to support people in distress, including advertising crisis hotline numbers. Identify gaps in the continuum of crisis services, such as warm lines to reduce loneliness and isolation and access lines to connect people to local resources. Deliver suicide prevention training to all crisis service providers.
- Disseminate information on available crisis service resources to health and behavioral health care partners to ensure resources are included in safety plans developed through an alliance between providers and people at risk.
- Create memorandums of understanding between systems of care and community-based crisis services to provide follow-up for people transitioning out of care systems, including protocols for protecting the confidentiality of people at risk. Health and behavioral health care systems should have protocols in place for obtaining consent for follow-up care from people at risk. To coordinate efforts, document clear methods of communication between crisis service providers and other systems, such as community corrections, child welfare, and veterans' services.

Action to Implement State Strategies

Goal 8:	Increase detection and screening to connect people to services based on suicide risk
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of increasing detection and screening to connect people to services based on suicide risk.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 3.
Expand resources to support health care practitioners and providers increase access and linkage to behavioral health services for people identified as needing such services.	By July 1, 2022, the State, in consultation with private and public partners, should create incentives to expand the use of Collaborative Care in health care systems. Options may include expanding the scopes of practice for physician assistants and nurse practitioners specifically trained in suicide prevention risk assessment, management, and referral, creating guidance and reducing barriers for billing health plans for services, and reducing documentation burden.
Adopt the Zero Suicide Initiative within health and behavioral health care systems.	By January 1, 2023, the State, in consultation with private and public partners, should form a task force to make recommendations for implementing the Zero Suicide Initiative framework into public and private health and behavioral health care systems across the state, including identifying state funds that may be needed to build capacity for technical assistance and training. As part of this effort, the department should partner with California health systems currently implementing the Zero Suicide Initiative, such as Kaiser Permanente.
Increase standardized training in best practices in suicide risk assessment and management to health care practitioners and providers and enhance uniform suicide risk assessment and management in health care settings, in alignment with Joint Commission	<p>By July 1, 2023, the State, in consultation with private and public partners, should develop a strategy for delivering training in best practices for suicide risk assessment and management to all health care practitioners and providers. Because health care practitioners and providers are at increased risk for suicide themselves, trainings should include a component on best practices for provider wellness, including methods of reducing burn-out, compassion fatigue, and vicarious trauma.</p> <p>By December 31, 2022, the Office of Suicide Prevention should disseminate guidance for screening for suicide risk for at-risk</p>

Goal 8:	Increase detection and screening to connect people to services based on suicide risk
State Strategy	Action to Implement Strategy
guidelines and the Zero Suicide Initiative.	groups, including people exposed to physical and sexual abuse, victims of domestic or other interpersonal violence, families and youth in the child welfare system, and people in detention settings or on probation or parole supervision.
Identify opportunities for technology advancements in health care systems to advance suicide risk assessment and management application, use, and investment.	By July 1, 2023, the Office of Suicide Prevention should form a task force to develop and disseminate guidance for the use of technology to support suicide risk assessment and management in high-risk settings, including health care systems. This effort also should explore the potential to use administrative data to detect and monitor suicide risk when screening is not feasible. For example, school administrative data indicating risk might include absences, excessive tardiness, and significant changes in academic performance and behavior in school.

Goal 9:	Deliver a continuum of crisis services within and across counties
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of delivering a continuum of crisis services within and across counties.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 3.
Develop and implement a strategy to coordinate the delivery of crisis services, including an assessment of current crisis services infrastructure and private and public funding for services.	By July 1, 2022, the State, with leadership from the Department of Health Care Services and private and public partners, should form a task force to develop a strategy for evaluating crisis services and the extent to which crisis services prevent suicidal behavior and make recommendations for standardizing crisis service delivery systems across the state based on findings, including addressing training and capacity barriers. The evaluation plan should be implemented by July 1, 2023.

Goal 9:	Deliver a continuum of crisis services within and across counties
State Strategy	Action to Implement Strategy
	<p>As part of this effort, the State should assess current capacity for training and technical assistance needed to systematically improve crisis services statewide. The department should explore implementing the Crisis Now Model across California. The department also should develop a process to monitor quality assurance and quality control of crisis services, including how the state will track data, targets, and measures regularly and report to the public. After assessing need and identifying private and public funding sources, the department should make recommendations to the Governor and Legislature any additional resources required to ensure the crisis services network is sufficiently funded. The department could use a tool, such as the Crisis Resource Need Calculator, for its assessment.</p> <p>By December 31, 2022, the Office of Suicide Prevention should develop and disseminate guidance for planning and coordinating crisis services for schools, colleges, and universities to prevent suicidal behavior among students. The guidance should include information about how schools could formally connect to crisis services and supports in the community.</p> <p>By December 31, 2022, the Office of Suicide Prevention should develop and disseminate guidance for integrating best practices in suicide prevention in crisis intervention training and co-responder models when law enforcement and mental health providers respond to behavioral health crises. The best practices should include assessment and referral to services based on suicide risk and creating safety by reducing access to lethal means.</p>
Develop a state accreditation program for suicide and crisis hotlines operated in the state, including standards for training and core	By December 31, 2021, the Office of Suicide Prevention should develop a strategy for collecting crisis services data and monitoring the quality, timeliness, and effectiveness of services to reduce suicidal behavior.

Goal 9:	Deliver a continuum of crisis services within and across counties
State Strategy	Action to Implement Strategy
competencies for call responders, protocols for performance and quality assurance monitoring, and procedures for making referrals to services, including emergency services.	As part of this effort, the office should develop standards for suicide prevention hotlines and centers, including training for hotline staff and performance targets. One option would be adopting minimum standards set by an accrediting organization, such as the American Association of Suicidology or the National Suicide Prevention Lifeline. The office should identify incentives for accreditation, such as requiring accreditation as a condition for state funding.

Strategic Aim 4: Advance suicide-specific services and supports

Goal 10: Deliver best practices in care targeting suicide risk

Strategic Aim 4: Advance suicide-specific services and supports

Short-Term Target

- By 2025, 30 percent of practitioners and providers receive standardized training in best practices in suicide risk assessment, management, and treatment

Long-Term Outcome

- Decrease in suicidal behavior

Key Action Partners	
<ul style="list-style-type: none">• Survivors of suicide attempt and loss• Health, public health, and behavioral health leaders and providers• Indigenous and traditional healers	<ul style="list-style-type: none">• Crisis services providers• Suicide prevention organizations• Hospital representatives• Parents and caregivers

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of delivering best practices in care targeting suicide risk.
- Create a process to certify providers trained in delivering best practices in treating people at risk for suicide. Such processes could include minimum education, training, and continuing education requirements, and should include a review and approval process prior to certification.
- Create a strategy to increase workforce capacity for providers to deliver suicide-specific care.

Local Strategies to Support Goal

- Expand the use of telehealth and telemedicine providers with training in best practices for suicide-specific treatment - especially in rural communities - to enhance timely access to care targeting suicide risk.

- Disseminate information to promote safety planning by prompting health care practitioners and providers to record safety plans in electronic medical record systems and by making plans accessible to people via commonly used portals.²⁷
- Create a local online network of providers delivering suicide-specific treatment, including public information about insurance eligibility and criteria for new clients.
- Partner with health care systems and providers to advance delivery of services and supports to caregivers and family members of people transitioning from health care settings following services for suicidal behavior. Prioritize safety and address service gaps. People at risk should be key decision-makers in defining support networks and the role each member of the network plays in creating safety and recovery.
- Disseminate information to caregivers and family members on how they can support a person at risk by being a resource identified by the person in safety planning; reduce environmental safety risks by promoting means safety, especially at home; and help manage harmful behaviors stemming from underlying health and behavioral health needs, such as escalating alcohol or drug use.

Goal 11: Ensure continuity of care and follow-up after suicide-related services

Strategic Aim 4: Advance suicide-specific services and supports

Short-Term Targets

- By 2025, all people prior to being discharged from emergency departments and hospital settings after receiving suicide-related services are connected to a provider to develop scheduled follow-up contacts over a 12-month period or more, as needed
- By 2025, 20 percent of those discharged from an emergency department or hospital setting after receiving suicide-related services receive outpatient services within 48 hours following discharge

Long-Term Outcome

- Reduce subsequent suicidal behavior among people discharged from emergency departments and hospital settings after suicide-related services

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • Health, public health, and behavioral health leaders and providers • Indigenous and traditional healers • Parents and caregivers 	<ul style="list-style-type: none"> • Crisis services providers • Suicide prevention organizations • Hospital representatives • Youth leaders

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of ensuring continuity of care and follow-up after suicide-related services.
- Develop best practice guidance for ensuring continuity of care following discharge after suicide-related services in emergency departments and hospital settings, including routine and standardized follow-up cards and notes.
- Establish a program to deliver training on lethal means restriction counseling to health care practitioners and providers and distribute gun and medication lock boxes and locks to hospitals, with prioritized distribution to people being discharged following a suicide attempt.

Local Strategies to Support Goal

- Increase the use of electronic health records to document a person's safe transition to another provider and ensure life-saving information is transmitted, while protecting privacy.
- Facilitate safe and timely care transitions through linkages to outpatient behavioral health providers, crisis services, safety planning or crisis response planning, and by reducing access to lethal means.
- Disseminate the *Caring for Adult Patients with Suicide Risk: A Consensus Guide for Emergency Departments* found at http://www.sprc.org/sites/default/files/EDGuide_full.pdf, to emergency department administrators, along with the *Quick Guide for Clinicians* found at http://www.sprc.org/sites/default/files/EDGuide_quickversion.pdf, to increase awareness of safe discharge practices for people seen for suicide-related services.

- Train health care practitioners and providers to deliver lethal means counseling to family members and caregivers supporting discharge from a health care setting for suicidal behavior.
- Disseminate information on lethal means counseling to health care practitioners and providers across hospital settings. Prioritize providers who predominately serve at risk-groups or providers in high-risk settings, such as emergency departments. Promote free online training, such as Counseling on Access to Lethal Means available at <https://training.sprc.org/>, and the use of online toolkits, such as <https://health.ucdavis.edu/what-you-can-do/>.
- Create uniform policies and procedures for safely transitioning people or students back into the workforce and home or school following a suicide attempt, suicide loss, or hospitalization for a behavioral health crisis.
- Create uniform policies and protocols to support health care practitioners and providers in revising or creating a safety plan with a person at risk, establishing a connection between the person and the new care provider, ensuring timely delivery of information to the new care provider, and following up within 24 to 48 hours of the transition. Create memorandums of understanding with local crisis service providers to establish relationships with people prior to discharge and follow-up after release.
- Create uniform protocols for counseling people discharged from emergency departments and hospitals after receiving suicide-related services on restricting access to lethal means. Families and caregivers should be included in such counseling.

Goal 12: Expand support services following a suicide loss

Strategic Aim 4: Advance suicide-specific services and supports

Short-Term Target

- By 2025, all counties have written policies and procedures for coordinated, timely, and respectful responses by service providers following a suicide loss, including formal agreements with local coroners and medical examiners to support the initiation of services

Long-Term Outcome

- Increase in the number of bereavement services specifically designed to meet the needs of suicide loss survivors

Key Action Partners	
<ul style="list-style-type: none"> • Survivors of suicide attempt and loss • First responders • Health, public health, and behavioral health leaders and providers • Parents and caregivers • Youth leaders • Coroners and medical examiners 	<ul style="list-style-type: none"> • Hospital representatives • School administrators • Faith-based community leaders • Chaplains • Mortuaries • Suicide prevention organizations

State Strategies to Support Goal

- Create a research and policy agenda to advance the goal of delivering support following a suicide loss.
- Assess and expand effective resources available to suicide loss survivors and develop capacity statewide to deliver appropriate and respectful services following a suicide loss.

Local Strategies to Support Goal

- Develop an integrated postvention services plan to guide delivery of best practices following a suicide loss. The plan should tailor strategies to settings and cultures, including schools, workplaces, faith communities, hospitals and health care settings, tribal communities, and correctional facilities. The plan should identify a lead agency or organization responsible for ensuring adequate capacity, training, and effectiveness in the delivery of activities that support survivors, service providers, and community members after a suicide loss. Enter into agreements to clearly define roles and procedures for a coordinated response.
- Develop an online bereavement toolkit with community-specific resources and partner with hospitals, first responders, funeral directors, faith-based communities, and coroners and medical examiners to distribute print copies or web links.
- Deliver training to first responders, crisis service providers, and access line responders on best practices for supporting suicide loss survivors, including understanding unique needs and available resources.
- Create local suicide bereavement support programs or expand capacity and sustainability of existing programs using *Pathways to Purpose and Hope* found at

<https://emmresourcecenter.org/resources/pathways-purpose-and-hope-guide-creating-sustainable-suicide-bereavement-support-program>.

- Expand support services designed and facilitated by survivors of suicide loss. Train survivors of suicide loss to speak safely and effectively about their loss and create a local speakers bureau to give a forum for survivors to deliver suicide prevention messaging to the public. Provide training for suicide loss survivor service facilitators, and opportunities for service facilitators to support each other, including group debrief sessions.
- Enter into memorandums of understanding with coroners and medical examiners to establish coordinated, timely, and respectful responses following a suicide loss, and establish policies and protocols to govern activities in the event of a suicide. Components should include how information is shared, and with whom, and how the privacy of the family is respected, including a process for determining how and when to reach out to family members with resources and support.

Action to Implement State Strategies

Goal 10:	Deliver best practices in care targeting suicide risk
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of delivering best practices in care targeting suicide risk.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 4.
Create a process to certify providers trained in delivering best practices in treating people at risk for suicide. Such processes could include minimum education, training, and continuing education requirements, and should include a review and approval process prior to certification.	By July 1, 2023, the State, in consultation with private and public partners, should create incentives for behavioral health licensing entities to develop a certification specific to providers who can deliver best practices in treatment targeting suicide risk and to develop a database of all certified practitioners and providers that is accessible to the public. California's mental health licensing entities include the Medical Board, the Board of Psychology, and the Board of Behavioral Sciences.
Create a strategy to increase workforce capacity for providers to deliver suicide-specific care.	By December 31, 2022, the Office of Suicide Prevention should develop an online resource center to support the continuing education of best practices in suicide prevention interventions and therapies for health and behavioral health care providers. By December 31, 2024, the State, in consultation with private and public partners, should require education and training in best practice therapies targeting suicide risk in all medical and clinical education training curriculum.

Goal 11:	Ensure continuity of care and follow-up after suicide-related services
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goals	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a

Goal 11:	Ensure continuity of care and follow-up after suicide-related services
State Strategy	Action to Implement Strategy
of ensuring continuity of care and follow-up after suicide-related services.	research and policy agenda to advance the goals outlined in Strategic Aim 4.
Develop best practice guidance for ensuring continuity of care following discharge after suicide-related services in emergency departments and hospital settings, including routine and standardized follow-up cards and notes.	<p>By July 1, 2023, the State, in consultation with private and public partners, should require all hospitals and emergency departments to develop policies and protocols for delivering counseling on lethal means restriction; distributing means safety products, such as lock boxes for guns or medications; and sending follow-up messages to people discharged after receiving services for a suicide attempt.</p> <p>Protocols and practices must include provisions detailing how informed consent will be obtained and how follow-up care will reflect a collaborative, transparent approach with the person at-risk to prioritize outpatient care. Protocols and procedures must include brief interventions involving best practices in safety planning and lethal means counseling. Follow-up care must be linguistically and culturally respectful. Protocols and practices should include methods for tracking linkages to referrals to services, when possible.</p> <p>By July 1, 2023, the Office of Suicide Prevention should form a task force to develop and disseminate best practice guidance and make recommendations for comprehensive aftercare for people discharged from hospital settings, including standardizing a process for delivering follow-up, establishing care linkages prior to discharge, and ensuring ongoing monitoring and support.</p> <p>Guidance should highlight the role of California’s suicide prevention hotlines and centers in providing proactive services by establishing a connection with suicide attempt survivors prior to discharge and conducting routine follow-up to ensure connections to services. Guidance should include opportunities to increase “rapid referrals” and should identify incentives for health care practitioners and providers. These referrals involve people who either are</p>

Goal 11:	Ensure continuity of care and follow-up after suicide-related services
State Strategy	Action to Implement Strategy
	<p>being treated in an emergency department or are approaching hospital discharge, and the goal is to connect them from inpatient care to outpatient services within 24 to 48 hours after discharge.</p> <p>By July 1, 2023, the State, in consultation with private and public partners, should create incentives for outpatient behavioral health care providers to enter into agreements with hospitals to accept referrals and develop a process for confirming timely outpatient appointments prior to discharge.</p> <p>By July 1, 2024, the Office of Suicide Prevention should partner with schools, universities, and colleges to identify challenges and opportunities for safely transitioning students back into schools after hospitalization for suicidal behavior and develop and disseminate best practice guidance.</p>
Establish a program to deliver training on lethal means restriction counseling to health care practitioners and providers and distribute gun and medication lock boxes and locks to hospitals, with prioritized distribution to people being discharged following a suicide attempt.	By July 1, 2023, the State, in consultation with private and public partners, should create a program to support training for health care practitioners and providers and hospitals in distributing means safety products, such as lock boxes for guns or medications, to people discharged after receiving services for a suicide attempt.

Goal 12:	Expand support services following a suicide loss
State Strategy	Action to Implement Strategy
Create a research and policy agenda to advance the goal of expanding support services following a suicide loss.	By December 31, 2021, the Office of Suicide Prevention should form a task force of subject matter experts to create a research and policy agenda to advance the goals outlined in Strategic Aim 4.

Goal 12:	Expand support services following a suicide loss
State Strategy	Action to Implement Strategy
<p>Assess and expand effective resources available to suicide loss survivors and develop capacity statewide to deliver appropriate and respectful services following a suicide loss.</p>	<p>By July 1, 2022, the Office of Suicide Prevention should develop and expand capacity for a statewide network of survivor support service providers across settings, including in schools, workplaces, health care, faith communities, tribal communities, and correctional facilities.</p> <p>By January 1, 2023, the Office of Suicide Prevention should form a task force to evaluate services delivered to people bereaved by suicide loss and identify gaps in services and disseminate findings.</p> <p>By July 1, 2024, the task force should make recommendations for implementing best practices in local team-based responses following a suicide loss in a community or specific setting, including how to manage privacy and information and data sharing among members of the team.</p> <p>By July 1, 2024, the task force should develop guidance for coroners, medical examiners, and law enforcement for supporting people bereaved by suicide, including methods for reducing stigma and shame and responding to cultural differences following a suicide loss.</p>

Plan Development

The California Legislature passed Assembly Bill 114 (Chapter 38, Statutes of 2017), mandating that the Mental Health Services Oversight and Accountability Commission develop a statewide strategic suicide prevention plan. The Commission began the work in early 2018 by forming a Suicide Prevention Subcommittee, which included Commissioners Tina Wooton (Chair), Khatera Tamplen, and Mara Madrigal-Weiss.

Community Engagement and Site Visits

The Commission organized a series of meetings and events to identify challenges in suicide prevention and opportunities for improvement. These opportunities were designed to engage public discussion and ensure that statewide planning reflected California's unique cultural, ethnic, linguistic, and economic diversity. The meetings were public and sought to incorporate a broad range of perspectives to support the development of shared knowledge to advance strategic planning. Please visit www.mhsoac.ca.gov for a full list of community engagement activities and summaries from events.

The Subcommittee held meetings in Fresno, Sacramento, San Diego, and Shasta counties to hear presentations on local suicide prevention initiatives and explore with community members challenges and opportunities for suicide prevention. Several common priority areas emerged from these meetings: the need for early identification of suicide risk and methods for reducing isolation, increasing access to appropriate services, and opportunities for leveraging partnerships to build capacity. The Commission held two public hearings to explore with suicide loss and attempt survivors, providers, researchers, and other subject matter experts' recommendations for closing gaps in data collection, service delivery, and training and education.

Topic-specific workshops and forums were held to supplement presentations and discussions organized via subcommittee meetings and public hearings. Events were designed to gather perspectives from communities affected by suicide in ways where data and information are limited, including youth, first responders, and diverse communities. A common finding from these events was that suicide prevention efforts are most effective when they are culture-specific and include planning and delivery by people from the at-risk group. Project staff participated in the City of Los Angeles Mayor's Challenge to Prevent Suicide.²⁸ Project staff heard input from members of the California Department of Education's Student Mental Health Policy Workgroup, Indian Health Services, California Rural Indian Health Board, and many other organizations.

The Commission visited several sites in California to explore key opportunities for suicide prevention. Sites included the Rancheria Health Center and Counseling and Recovery

Engagement Center in Shasta County, UCSF Benioff Children's Hospital in Alameda County, and the Golden Gate Bridge.

Research and Subject Matter Expert Consultation

As part of its research for this report, project staff met with local and national leaders in suicide prevention. Staff worked with representatives of departments under the California Health and Human Services Agency as well as other state and local government and private partners. These included behavioral health, public health, law enforcement, and education officials as well as representatives of foundations, nonprofit organizations, the healthcare industry, and other businesses. Staff also engaged with national leaders from the American Foundation for Suicide Prevention, National Zero Suicide Initiative, National Action Alliance for Suicide Prevention, Suicide Prevention Resource Center, Centers for Disease Control, United States Substance Abuse and Mental Health Services Administration, and Suicide Awareness Voices of Education. Staff participated in a national convening of behavioral health and suicide prevention experts and attended a training on the Zero Suicide Initiative.

Finally, the Commission conducted a critical review of the latest research on suicide prevention best practices, along with other information gathered through local, national, and international efforts. Commission staff consulted national and global frameworks for preventing suicide, including, but not limited to:

- The 2012 National Strategy for Suicide Prevention, developed by the U.S. Surgeon General and the National Action Alliance for Suicide Prevention
- Public Health Action for the Prevention of Suicide: A Framework (2012) and Preventing Suicide: A Global Initiative (2014) by the World Health Organization
- Preventing Suicide: A Technical Package of Policy, Programs, and Practices by the Centers for Disease Control and Prevention and the Substance Abuse and Mental Health Services Administration

The Commission contracted with content experts at Stanford University to provide technical guidance on research and best practices in suicidology and public health strategy. Suicidologist Dr. Rebecca Bernert led the team of technical advisors, which included Drs. Keith Humphreys and Shashank V. Joshi.

Previous Suicide Prevention Plan

Development of the current suicide prevention plan included reviewing the previous suicide prevention plan approved in 2008. In September 2006, Governor Arnold Schwarzenegger directed the former Department of Mental Health to develop a statewide strategic suicide prevention plan. The plan was approved by the Governor's Office on June 30, 2008. Many of the actions recommended were not fully implemented. The current plan retains much of what was proposed, with updated best practices in means restriction, health care, and data monitoring and evaluation. Key advancements directed by the previous plan – at least partially fulfilled - are briefing highlighted below.

Leadership

The 2008 plan called for a dedicated state office to provide coordination and collaboration across the state. The Office of Suicide Prevention was established by the Department of Mental Health but was transferred and reorganized into the Suicide Prevention Program after the department was closed in 2012.²⁹ The program is currently at the Department of Health Care Services.³⁰ Core functions of the Office of Suicide Prevention, such as convening regional meetings, disseminating of resources to local county suicide prevention liaisons, and coordinating suicide prevention activities to advance the goals under the plan have since ended.

Guidance for Policy and Practice

Local suicide prevention activities have expanded since 2008, largely through funding with Mental Health Services Act dollars. A portion of the funding is directed toward the prevention of the consequences of unmet mental health needs, including suicide. County behavioral health departments use this funding by reducing risk factors for mental illness through “prevention programs” and “early intervention programs,” and by initiating suicide prevention programs that specifically prevent suicide as a consequence of mental illness.³¹ Departments spent over \$13 million during FY 2016-17 on suicide prevention activities, such as suicide prevention hotlines, gatekeeper training, screening for older adults, and services supporting suicide loss survivors, among many others.³²

Several counties in California have suicide prevention plans and local task forces or collaboratives with multi-disciplinary partners working together to prevent suicide. Counties that have created local plans include Contra Costa, Fresno, Kings, San Diego, San Mateo, Santa Clara, Solano, Tulare, and Tuolumne. Counties that have local collaboratives include Contra Costa, Fresno, Kings, Los Angeles, Napa, Nevada, San Diego, San Mateo, Shasta, Solano, Tulare, Tuolumne, and Ventura. Other counties, such as Marin, Santa Cruz, and Stanislaus Counties are in the planning phase. For example, Stanislaus County was approved to use Mental Health Services Act Innovation funding to use collective impact principles to develop a local suicide prevention plan but does not have a plan in place at this time.³³

California public schools with students in grades seven through 12 are now required to develop a suicide prevention policy, known as the Pupil Suicide Prevention Policy. The policy must be created in consultation with school and community stakeholders, school-employed mental health professionals, and suicide prevention experts, and must include procedures related to suicide prevention, intervention, and postvention. All policies were to be in place by the 2017-18 school year. A review conducted in 2018 by the Trevor Project found that 86 percent of schools that are legally required to have plans have them in place, leaving approximately 69 schools without plans.³⁴

Training

One of the goals of the 2008 plan was to develop and implement training and workforce enhancements to prevent suicide. Legislation was passed in 2017 requiring licensed psychologists to receive no less than six hours of training in suicide risk assessment and intervention by 2020.³⁵ Additional legislation was passed in 2018 to extend this requirement to mental health professionals licensed by the Board of Behavioral Sciences.³⁶ In addition to increased training for clinicians, the Legislature allocated \$1.7 million for one-time general funding for online suicide prevention training for all public middle and high school students and staff in California.³⁷ Despite these critical advancements, there still remains a need for standardized training guided by best practices.

Technical Assistance

The 2008 plan outlined the need for technical assistance, such as establishing regional learning collaboratives, training guidance, an online clearinghouse, and support for local suicide prevention efforts. The Commission approved one-time Mental Health Services Act funding of \$40 million over four years for statewide infrastructure, such as a clearinghouse of best practices to assist in training and technical assistance efforts, as well as a suicide hotline system, which would benefit all counties.³⁸ That investment resulted in several initiatives administered by the California Mental Health Services Authority – some of which are still operational.³⁹ These initiatives created regional networks focused on collaboration and development of best practices, delivered suicide prevention training, developed social media marketing campaigns, and partnered with crisis centers to expand cultural and linguistic competent outreach, technology capacity to chat and text functions, and improved data collection.⁴⁰ Highlighted in this section are the Know the Signs Campaign, the Directing Change program and film contest, and the California Suicide Prevention Network.

The Know the Signs Campaign is a social marketing initiative to educate Californians on how to recognize the warning signs of suicide, how to talk to someone in crisis, and how to access services.⁴¹ The campaign also works with members of the media to promote consistency with national recommendations for reporting suicides in the media. Directing Change is a program and film contest in California designed to engage students in creating films to promote positive conversations about mental health and suicide prevention.⁴² Lastly, the California Suicide Prevention Network was established using the initial funding to centralize statewide suicide

prevention activities, reduce stigma associated with suicide, and increase access to care for people at risk of suicide. The network produced common metrics for evaluating suicide prevention hotlines: demographic data of callers, reason for call, call volume, and suicide risk of caller.

Suicide Hotline Assessment

One next step identified in the 2008 plan was to assess the current status of coverage and accreditation for suicide prevention hotlines.⁴³ The Department of Health Care Services was directed in 2016 to conduct a comprehensive assessment of suicide hotlines and recommend funding strategies to ensure hotlines have adequate resources to meet demand.⁴⁴ The department produced a report that documented the structure, capacity, and funding of current suicide hotlines accredited by the American Association of Suicidology across the state.⁴⁵ The report highlighted the demand for a statewide suicide hotline system but said a lack of data prevented the department from determining funding needs to meet demand.⁴⁶ Currently, \$4.3 million per year of Mental Health Services Act funding is allocated to support suicide prevention statewide through a suicide hotline provider, in addition to local and private funding sources.⁴⁷

Public Review

The draft statewide strategic suicide prevention plan was first released for public comment on July 2, 2019. The Subcommittee received written and verbal comments before the plan was submitted to the Commission for consideration.

Plan Note

This plan does not include physician-assisted dying, which is sometimes referred to as assisted suicide. In California, the End of Life Option Act allows qualified adults with a terminal illness to request aid-in-dying drugs from their physician.⁴⁸

Suicidal Behavior: Definitions, Theory, and Key Concepts for Prevention

Suicidal behaviors exist on a broad continuum of risk, and include desire to die and suicidal ideation, suicide attempt planning, suicide attempts, and death by suicide. The Centers for Disease Control and Prevention uses the term self-directed violence to describe a range of violent behaviors that can be fatal or non-fatal, suicidal or non-suicidal.⁴⁹ Suicide is defined as “death caused by self-directed injurious behavior with any intent to die as a result of the behavior.”⁵⁰ For the purposes of this document, non-fatal, suicidal self-directed violence is referred to as “suicidal behavior” to represent the full continuum of risk.

Non-Suicidal and Suicidal Self-Directed Violence Definitions

Self-directed violence is behavior that is self-directed and deliberately results in injury or the potential for injury to oneself.⁵¹ Behavior can be non-suicidal or suicidal.

Non-suicidal self-directed violence is behavior that is self-directed and deliberately results in injury or the potential for injury to oneself, with no evidence - implicit or explicit - of suicidal intent.

Suicidal self-directed violence is behavior that is self-directed and deliberately results in injury or the potential for injury to oneself, with evidence – implicit or explicit - of suicidal intent. Suicidal self-directed violence includes:

- **Suicide attempt**, a non-fatal, self-directed potentially injurious behavior with any intent to die as a result of the behavior. A suicide attempt may or may not result in injury.
- **Interrupted or aborted suicide attempt**, an effort to injure oneself that is stopped by the person attempting self-harm, or by another individual prior to fatal injury. This can occur at any point during the act, such as after the initial thought or after the onset of behavior.
- **Preparatory acts** or preparation toward making a suicide attempt, taken before potential for harm has begun. This can include any action beyond a verbalization or thought, such as purchasing a gun or preparing for one’s death by suicide by giving away belongings.

Suicidal behavior also can include suicidal ideation, defined as having the desire to die, or thinking about engaging in behaviors to die.⁵² Suicidal ideation can be passive or active.⁵³ If active, suicidal ideation can be nonspecific, can include a method but no intent or plan, can include a method and intent but no plan, and can include method, intent, and plan.⁵⁴ For the purposes of this document suicidal ideation is referred to as suicidal behavior, unless specified

Suicidal Ideation Definitions and Screening

Five levels of suicidal ideation – increasing in severity - are outlined within the Columbia-Suicide Severity Rating Scale:⁵⁵

Suicidal Desire

Person has a wish to be dead or not alive, or a wish to fall asleep and not wake up.

Suicidal Ideation (Thoughts) – without thoughts of method

Nonspecific thoughts about suicide or wanting to end one's life, without thoughts of a method for an attempt. Example: *Life is not worth living.*

Suicidal Ideation: Includes method - no intent or plan

No specific plan with time, place, or method details worked out. Example: *I've thought about driving off the road or overdosing, but never of acting on the thought.*

Suicidal Ideation: Includes method and some intent - but no plan

Thoughts of an attempt method, with some intent to act. Example: *I've thought about driving off the road and have thought about acting on it when feeling at my worst.*

Suicidal Ideation: Includes method, intent, and plan

Thoughts of attempting suicide with details of a plan and some intent to carry it out.

Example: *I've started to work out plans for how to overdose and intend to carry it out.*

The Columbia-Suicide Severity Rating Scale uses the following questions to screen for severity of suicidal ideation and is used to support decisions for services and referral based on risk:

1. Have you wished you were dead or wished you could go to sleep and not wake up?
2. Have you had any thoughts of suicide?
3. Have you been thinking about how you might do this? *For example, "I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it....and I would never go through with it."*
4. When you had these thoughts, did you have some intention of acting on them? *As opposed to "I have the thoughts but I definitely will not do anything about them."*
5. Have you started to work out or worked out the details of how to attempt suicide? Do you intend to carry out this plan?

See www.cssrs.columbia.edu for downloadable measures designed for select settings and groups.

Assessing for Suicide Risk

Suicidal ideation and risk level may vary according to the *intensity, duration, pervasiveness, the controllability of symptoms, reasons for living, and history of past suicide attempts or non-suicidal self-injury*.⁵⁶ As a result, the Columbia-Suicide Severity Rating Scale and other assessment measures prioritize evaluation of the intensity of suicidal ideation, such as asking about duration, controllability, deterrents, reasons for the thoughts, as well as evaluation of suicidal behavior, such as history of suicide attempt, interrupted or aborted attempt, preparatory behaviors, and intentional self-harm without desire or intent to die.⁵⁷ Such factors in suicide risk assessment are discussed in greater detail in subsequent sections that review best practices in collaborative assessment and management of suicide risk. These suicide risk assessments use a collaborative and transparent approach to assessing for suicide risk and to support delivery of additional services, referral, or safety planning.

Suicide Theory

Suicide is a complex public health challenge involving many biological, psychological, social, and cultural determinants.⁵⁸ Several theories about why people die by suicide seek to explain how multiple factors may increase risk in the context of profound emotional suffering. According to the Interpersonal Theory for Suicide, three components must align to predict risk for suicide or a serious suicide attempt: thwarted belongingness, perceived burdensomeness, and acquired capability for lethal self-injury.⁵⁹

Thwarted Belongingness and Perceived Burdensomeness

The Interpersonal Theory for Suicide includes two components of suicidal desire and depression: “thwarted belongingness” and “perceived burdensomeness.”⁶⁰ Thwarted belongingness is described as a state of “unmet need to belong.”⁶¹ Both the theory and extensive research indicate that people have a fundamental need to belong and that, when that need is thwarted, it increases risk.⁶² A sense of belonging can increase during times of national celebration and in times of national crisis, such as during wartime. One illustration of this involved the change in the national daily suicide rate following the attacks on September 11, 2001.⁶³ In the year following the attacks, suicide rates in entire U.S. communities showed an unprecedented decrease – but only on that day, not in the period before or after.⁶⁴ Similar findings are observed in times of national celebration.⁶⁵ Perceived burdensomeness is the false belief that “my death is worth more than my life.”⁶⁶ Unemployment, health problems, and incarceration are examples of situations in which a person may feel like they are a burden to others. This finding aligns with empirical research indicating these situations increases risk for suicide.⁶⁷

Acquired Capability

Although the components described above are components of depression and reflect suicidal desire, which are modifiable and amenable to services, the Theory proposes that these factors are not on their own predictive of risk. Indeed, most people with depression do not go on to die by suicide. The Theory instead proposes that people are most at risk when these components are present in combination with an acquired capability for self-injury, or “the ability to engage in suicidal behaviors acquired through life experiences that habituate pain tolerance and fearlessness about death.”⁶⁸ Such experiences may include exposure to physical pain, violence, and provocative life experiences, such as childhood trauma, witnessing a traumatic event, suffering from a chronic medical illness, or engaging in self-directed violence.⁶⁹ Indirect exposure to others’ pain and injury also may increase acquired capability, increasing risk among groups such as veterans, physicians, nurses, and first responders.⁷⁰

Means Matter

While reducing access to lethal means is a central element in global and national suicide prevention plans, it remains poorly understood and underutilized for reducing suicide in California.⁷¹ Suicidal behavior is often method-specific, and a person’s choice of means is driven by multiple factors. These include the lethality, accessibility, and acceptability of the method.⁷² Eliminating or reducing access to a particular method during a suicidal crisis creates lifesaving time and opportunity for intervention.⁷³ This is important because crises tend to be transient, and characterized by extreme ambivalence about the wish to die or stay alive.⁷⁴ Research shows that when a person’s attempt is thwarted, he or she does not go on to die by suicide at other locations, times, or by other methods.⁷⁵ As such, the placement of time between suicidal thoughts and a person’s ability to obtain lethal means for an attempt represent a practical, lifesaving approach to prevent suicide.⁷⁶

Gun access is a significant consideration in suicide prevention because the majority of people who die by suicide use a firearm.⁷⁷ While drug overdose is the most common method of suicide attempt, firearms are the most lethal.⁷⁸ Only about 15 percent of people who attempt suicide with a firearm will survive.⁷⁹ Using a highly lethal method of dying by suicide does not necessarily indicate a stronger desire to die.⁸⁰ Death by suicide is the result of many contributing factors, including choice of means, preexisting health or behavioral health needs, and the amount of time lapsed before rescue or medical intervention, among others. Lethality of means increases with age and escalates with the number of suicide attempts.⁸¹

Inherent Challenges and Emerging Innovations

Due to the nature of suicide, there are several inherent barriers to preventing it, making the implementation of comprehensive suicide prevention efforts challenging.⁸² These challenges are not immutable, but overcoming them will require a concerted effort.

Mental Health and Suicide Stigma

Harmful myths and stigma may discourage people from seeking help, prevent people from disclosing suicide risk, and hinder intervention and innovative in prevention and access to services. If left unaddressed, stigma can prevent multidisciplinary coordination across public and private industry partners, settings, and philosophies, and reduce the likelihood that suicide prevention will be included in public health strategies.⁸³ For example, though the majority of deaths by firearm occurs by suicide, suicide prevention and lethal means restriction are rarely discussed in gun safety campaigns and initiatives that promote safe gun storage.⁸⁴ Stigma also may affect public awareness of available services or effective practices to prevent suicide. Stigma likewise prevents people from seeking help for mental health needs. Stigma is also tied to disparities in seeking services for mental health needs and health access.⁸⁵ Men, for example, are more likely to receive mental health services in emergency departments because of perceived stigma associated with receiving mental health care. Understanding these disparities may help to identify targeted strategies for prevention and education training.

Disparities in Health Care Access

Suicide prevention services may be dependent on people at-risk seeking the services they need, which pose undue burden on people who may be in crisis. This challenge remains despite the effectiveness of screening protocols to guide triage and referral.⁸⁶ Services that specifically address suicide risk may be limited to select settings, such as a single community hospital, which limits the capacity to deliver integrated health care services across settings.⁸⁷ Variability in clinical practices may stymie the delivery of effective programs, while rural communities may experience shortages in services, especially for people with complex needs.⁸⁸

While psychosocial treatments for suicidal behaviors are effective, access to specialized care providers trained in such methods may limit wide-scale access, use, and adoption.⁸⁹ Insurance coverage may create barriers if people are unable to see specialists. Non-medical settings, such as the workplace or community centers, may be underutilized as opportunities to connect people with systems of care. These limitations may prevent services and effective approaches from being scaled statewide, or even within the same community.⁹⁰ Uniform guidelines for establishing visible and easily accessible pathways to access services has the potential to bridge this gap. Such guidelines could include centralized online resource hubs, provider referral networks with clearly described eligibility criteria, and standard protocols for best practices in transferring mental health emergency calls answered by 911 dispatchers to mobile crisis units or teams.

Missed Detection

Despite detection efforts, people at-risk for suicide may not be identified and receive the services they need when they need them.⁹¹ This challenge may be addressed by suicide prevention efforts integrated into entire systems intended to ensure people at-risk do not fall through gaps. Nationally, as of July 1, 2019, all people seen in medical settings for a primary diagnosis or primary complaint of a behavioral health need, including those seen in emergency departments as well as outpatient and inpatient settings, will be screened for suicide risk.⁹²

Other major suicide prevention initiatives in healthcare are underway. The Zero Suicide Initiative is an international movement toward systems transformation dedicated to preventing suicide within health care systems, with available free toolkits and training programs.⁹³ The majority of those who die by suicide interact with their doctor and health care system in the weeks and months prior to death.⁹⁴ The initiative promotes a system of continuous quality improvement in which health and behavioral health care providers develop policies and implement practices known to prevent suicide.⁹⁵ The potential to eliminate suicide when best practices are used and those risk are uniformly connected to evidence-based services is demonstrated through the Henry Ford Health System's Perfect Depression Care program, upon which the initiative is based.⁹⁶ Essential elements of the initiative are:

1. **Lead** systemwide culture change committed to reducing suicides
2. **Train** a competent, confident, and caring workforce
3. **Identify** people in care settings with suicide risk via comprehensive screenings
4. **Engage** all people at risk of suicide using a suicide care management plan
5. **Treat** suicidal thoughts and behaviors using evidence-based treatments
6. **Transition** individuals through care with warm hand-offs and supportive contacts
7. **Improve** policies and procedures through continuous quality improvement

Recent innovations in technology also offer hope for improving the detection of suicide risk, presenting opportunities for greater precision as well as increased screening sensitivity and better triage of people into services.⁹⁷ Machine learning is a form of Artificial Intelligence that enables a computer to learn patterns without prior programming and to devise complex algorithms to improve the accuracy of prediction.⁹⁸ Data routinely collected through electronic health records may be helpful in predicting future suicidal behavior.⁹⁹ An algorithm in one study of hospital-admission data – age, gender identity, zip code, medication, and diagnostic history, for example – was 84 percent accurate in predicting whether someone who was seen at the hospital for either non-suicidal self-injury or suicide attempt would attempt suicide in the following week.¹⁰⁰ The algorithm was 80 percent accurate in its prediction for a two-year period.¹⁰¹ Suicide prediction modeling is being developed for use in large healthcare systems, such as the U.S. Department of Veterans Affairs and Kaiser Permanente.¹⁰²

Machine learning also is being utilized by social media companies.¹⁰³ For years, Facebook users have had the ability to report posts by friends and family who they believed to be at risk for

suicide. In response to the posts, Facebook’s Community Operations team connects the flagged Facebook user with resources. Facebook has expanded its suicide prevention efforts by using machine learning to identify “suicidal expression” in posts by people at risk by monitoring phrases they use or comments from family and friends. Whether content is flagged by friends and family or by machine learning, the response is the same – a Community Operations team member reaches out to the person at risk, and, in emergencies, works with first responders.

Challenges in Terminology and Uniformity

Definitions for suicidal behavior are not uniform, and, likewise, there are no standards for suicide risk assessments, which affect risk detection, disclosure of risk, and reporting.¹⁰⁴ Despite calls for uniformity and national and state standards for screening, reporting, and data monitoring, there remain significant differences in how data are captured and how people are screened and referred to services.¹⁰⁵ Clinical practice guidelines for suicide prevention also reflect a lack of consensus, which may affect uniform procedures in risk assessment, triage, and training.¹⁰⁶ Differences in screening may hinder the ability to distinguish people at risk, preventing the delivery of effective programs and research of risk factors.¹⁰⁷ In response to these challenges, the Centers for Disease Control and Prevention created uniform guidelines to aid precision and comparability in the prevention and monitoring of suicidal behaviors.¹⁰⁸ Mandated screening and means restriction policies offer opportunities to aid detection given their universal use.¹⁰⁹

Barriers to Innovation

Despite advancements in suicide prevention, much is still unknown, and research exploring risk factors and treatments for suicidal behaviors remains a national and global priority. Specialists trained to conduct this research, however, are few relative to the need and priority. There is still much to understand about fundamental factors that contribute to risk for suicide and how risk changes over the lifespan, especially for specific groups.¹¹⁰ Risk factors change over time, and often are internal to each person. Visibility is key to detection of risk and intervention, as is the dissemination of information about how risk factors contribute to suicidal behavior and how those factors can be managed.¹¹¹ Finally, monitoring dynamic risk factors requires substantial and expensive infrastructure critical to building and sustaining effective suicide prevention initiatives.¹¹²

Research may be further hindered by funding and infrastructural barriers, and methodological, ethical, and safety challenges inherent to conducting epidemiological studies or research among those at high risk for suicide. Research on the effectiveness of interventions specifically targeting suicide risk is scarce. Until recently, people at risk for suicide were historically excluded from clinical drug trials due to safety concerns. This has limited the study of new treatments and is prioritized for drug safety and development, with now FDA-mandated assessment of suicide risk across all Central Nervous System drug trials.¹¹³

Suicidal Behavior in California

The following section describes suicidal behavior specific to California. It presents the state's suicide prevalence and rates based on the most recent data available. California's trends in suicide rates and suicidal behavior are aligned with national statistics, though some deviations are noted below. Trends in population and vulnerable group suicide rates are significantly affected by the method used for suicidal behavior; more lethal means, such as firearms, are involved in more suicide deaths.¹¹⁴

Suicide Data

In 2017, there were 4,323 Californians who lost their lives to suicide.¹¹⁵ California's age-adjusted¹ suicide rate is 10.7 per 100,000 people – one of the lowest rates among states – compared to the national rate of 14.0 per 100,000 people.¹¹⁶ California's relatively low suicide rate may be attributable to its policies regulating access and exposure to guns.¹¹⁷ In general, states with high rates of gun ownership tend to have higher rates of suicide and accidental death by firearm, whereas states with lower rates of gun ownership have lower suicide rates.¹¹⁸ While California's suicide rate is low compared to most other states, variability exists across counties. For example, Humboldt County has one of the highest suicide rates in California at 24.3 per 100,000 residents.¹¹⁹ Santa Clara County has the lowest suicide rate in California at 7.5 per 100,000 residents.¹²⁰ Variability in rates may be attributable to certain characteristics that increase risk for suicide, such as high gun ownership and less access to health care in rural communities.¹²¹

While rates are generally higher in rural Northern California counties, 2017 data show that a greater *number* of suicides claim the lives of residents in Southern California, specifically Los Angeles (21 percent of the state total suicides), Orange (10 percent of the state total suicides), Riverside (8 percent of the state total suicides), San Bernardino (6 percent of the state total suicides), and San Diego (5 percent of the state total suicides) counties, consistent with their population density.¹²² Half of all suicides in California in 2017 were reported in these five counties.¹²³ This concentration highlights the need for targeted, community-driven approaches and use of data to understand local and regional opportunities.

Suicide by Means

Firearm (37 percent), hanging and suffocation (32 percent), and poisoning, which includes overdose (16 percent), are the three most common ways people die by suicide in 2017, both nationally and in California.¹²⁴ Firearms were the leading cause of death by suicide, accounting for 37 percent of suicides in 2017.¹²⁵ Californians aged 30 and younger were more likely to die by hanging or suffocation while people older than 50 were more likely to die by firearm.¹²⁶ The

¹ Rates are adjusted using the 2000 US Standard Population weights and using 5 year age groupings for county and 10 year age groupings for the other variables. The age of the youngest suicide death is 10.

trend of younger people dying by suffocation is consistent with national trends.¹²⁷ Finally, 16 percent of Californians died by suicide from an overdose in 2017.¹²⁸

Suicide by Gender

In 2017, men died by suicide at a rate of more than three times higher than the rate of women in California.¹²⁹ This statistic is consistent with national data showing that men are nearly four times more likely to die by suicide than women.¹³⁰ This difference is largely explained by the use of more violent means among men.¹³¹ In other words, while attempt rates are higher for women, men are more likely to die as a result of an attempt because they use a firearm. Research consistently demonstrates that regardless of age group or culture, men are more likely to die by suicide and women are more likely to attempt suicide.¹³² Men dying by suicide at higher rates is consistent internationally, except for China where women – particularly young, rural residents - die by suicide at greater rates than men.¹³³

Suicide Rates by Age Groups

Risk of dying by suicide increases with age. In 2017, the suicide rate peaked at 14.5 per 100,000 for people between the ages 25 and 29, increased through middle-age, and was highest among Californians aged 85 and older (20.7 per 100,000 people).¹³⁴ This pattern is consistent with national trends. Californian men aged 85 and older had the highest suicide rate of any age group, at 45.1 per 100,000 people.¹³⁵ People in younger age groups attempt suicide at higher rates compared to older age groups but survive their attempt in part because of the selection of less lethal means for suicide.¹³⁶

Suicide Rates by Race/Ethnicity

Suicide rates in California are highest among whites (17.1 per 100,000 people) and Native Americans (15.6 per 100,000 people).¹³⁷ Native Hawaiian/Pacific Islander Californians had the next highest rate in 2017, at 14.1 per 100,000 people.¹³⁸ All other racial/ethnic group suicide rates were under 10 per 100,000 people.¹³⁹ This pattern is consistent with national trends, with white men accounting for nearly 70 percent of all suicide deaths in the U.S. in 2017.¹⁴⁰

Suicide by Military Members

There were 640 suicides by Californians aged 18 years and older who had served in the U.S. armed forces, accounting for 15.3 percent of all suicides in California in 2017.¹⁴¹ The majority of current and former military members who died by suicide were men (96.7 percent) and white (79 percent), and 43 percent were between the ages of 25 and 64 at the time of death.¹⁴² Additionally, 40 percent were between the ages of 65 and 84 at death.¹⁴³ The majority – 65.6 percent - of Californians who served in the Armed Forces and died by suicide in 2017 used a firearm.¹⁴⁴ Data showing that veterans are more likely than other at-risk groups to die by suicide using firearms highlights the need to consider as part of a prevention strategy the means by which different vulnerable groups die by suicide.¹⁴⁵

Suicide in Law Enforcement Custody

State and local law enforcement agencies are mandated to report the number of deaths in custody and arrest data, including death by suicide, to the California Department of Justice.¹⁴⁶ Custody settings include correctional housing, booking areas, holding cells, treatment units, and common areas, in addition to crime or arrest settings. Between 2005 and 2017, 922 people died by suicide in law enforcement custody.¹⁴⁷ The number of suicides in custody settings has decreased from a high of 83 in 2013 to 60 in 2017.¹⁴⁸ Most people who died by suicide in custody were male (93 percent) and were classified as white (49 percent), Hispanic (31 percent), or African American (11 percent).¹⁴⁹

Other Suicidal Behavior Data

In 2017, 18,153 Californians visited or were admitted to an emergency department for intentional self-harm.¹⁵⁰ Less is known about the prevalence of suicidal thoughts, because data may be limited to national or local self-report surveys. According to one survey, an average of 1,115,000 Californians over the age of 18 – about 3.8 percent of all adults – reported having serious thoughts of suicide in the past year.¹⁵¹ Another survey estimated that 19 percent of California 9th graders and 18 percent of California 11th graders seriously considered attempting suicide in the past year.¹⁵²

Data Limitations

There are many limitations to using current data to support suicide prevention efforts. Suicide is widely acknowledged as underreported as a manner of death on death certificates, both in the U.S. and internationally.¹⁵³ Manner of death includes suicide, homicide, accidental, or undetermined; cause of death refers to the circumstances of death, such as a gunshot wound. Coroners inquire into and determine the manner and cause of death when suicide is known or suspected.¹⁵⁴ After a death, a coroner or medical examiner follows procedures and protocols to investigate by documenting and evaluating the setting in which someone died; evaluating the body of the decedent; and evaluating medical, mental health, and social history.¹⁵⁵ Underreporting of suicide can occur because of inconsistent death classification.¹⁵⁶ While one coroner might label a death a suicide, another coroner confronted with the same circumstances might rule it “undetermined” or “accidental.” Cultural and religious beliefs, as well as stigma, also may influence accuracy of reporting and death records.¹⁵⁷

Several other barriers limit the use of suicide data for prevention efforts.¹⁵⁸ One is the inconsistent use by local jurisdictions of electronic reporting in centralized state databases, such as those maintained by the California Department of Public Health and the Office of Statewide Health Planning and Development.¹⁵⁹ Many death records remain in print form, which substantially delays reporting and real-time monitoring of suicide within and across

counties.¹⁶⁰ Further, bridge and railway suicide deaths are not reported in a unified manner by individual sites to a centralized reporting system, with information instead housed across multiple agencies, such as the California Department of Transportation (CalTrans), local transit districts, federal rail authorities, the California Highway Patrol, local sheriff-coroners, and other private entities.¹⁶¹ Compiling such data is crucial to evaluating public health risk and policy need, but a centralized reporting system is not currently in place.¹⁶²

Untimely data reporting and monitoring may also limit the ability of professionals to intervene when several suicides occur in proximity in place or time, known as a suicide cluster.¹⁶³ Inconsistent coding methods may compound difficulty when drawing comparisons between years, settings, or at-risk groups. In addition, data tends to be restricted to suicide deaths, despite critical opportunities for prevention in data associated with suicide attempts or “save data,” which describes a thwarted suicide attempt and subsequent connection to crisis services. For example, public data does not include how many people had repeat visits to the emergency department for suicidal behavior, data related to discharge or follow-up care outcomes, or first-time suicidal behavior not requiring triage services. These challenges highlight need to disseminate best practices statewide.

Risk and Protective Factors

Risk factors are characteristics that may make suicidal behavior more likely to occur, while protective factors are characteristics that make suicidal behavior less likely to occur.¹⁶⁴ Importantly, such factors often occur in the context of unmet health and behavioral health needs, interacting with other complex social, demographic, and situational dynamics. Factors that increase suicide risk, for example, are dangerous for people living with depression, while others can manage such factors. Some risk factors are modifiable, while others – such as history of suicidal behavior or demographic characteristics – are not. Effective suicide prevention efforts tend to target high-risk settings or risk and protective factors that can be modified, such as increasing screening and access to services depression and other behavioral health needs. Warning signs, by comparison, are behaviors that may indicate or signal acute risk for suicide, which may be similar to or distinct from risk factors.¹⁶⁵ See the next page for a list of risk and protective factors and warning signs.

Typically, risk can be elevated during times of acute or lasting transition, though the higher exposure is not limited to such periods. These transitions can include job loss, marital status changes, hospitalization, housing changes, and military service discharge or post-deployment. Risk appears to be additive – the more factors, the higher the risk – and it cuts across demographic, economic, social, and cultural boundaries. **Major risk factors for suicide are prior suicide attempt; substance abuse; mood disorder, such as depression; access to lethal means; and medical needs.**¹⁶⁶ Protective factors include the absence of risk factors and increased connectedness to community, culture, spiritual faith, and other factors that reduce risk, such as access to health care and social support and safely storing guns and medications. **Major protective factors for suicide are effective mental health care; connectedness to people, family, community, and social institutions; problem-solving skills; and contacts, such as postcards or letters, from service providers and caregivers.**¹⁶⁷

Cultural Considerations

Some risk and protective factors vary depending on the group targeted for suicide prevention efforts. For example, spirituality and religion are tied to reduced risk for suicidal behavior.¹⁶⁸ Spirituality and religion are deeply rooted in the culture, values, and norms of most ethnic groups.¹⁶⁹ Both can reinforce cultural transitions and strengthen cultural identity, protecting against risk.¹⁷⁰ Both may provide congregational opportunities to connect with community members, especially in times of stress, loss, and despair, reducing isolation and increasing resiliency and belonging. This can further mitigate risk by fostering hope and connection, promoting a sense of personal purpose or meaning, and teaching coping skills through spiritual practice.¹⁷¹

While religion is a protective factor for many communities, there are important differences among vulnerable groups. For example, religion may increase suicide risk among lesbian, gay,

bisexual, and transgender people.¹⁷² Adherence to religious doctrine that conflicts with sexual and gender identity can create confusion, distress, and isolation. This may be further compounded when people cannot seek support for their conflict and distress among members of their faith-based community.

Risk Factors

Suicide risk factors at the level of the person include:¹⁷³

- Prior suicide attempt(s)
- Suicidal thoughts with intent and planning (especially intense, pervasive, difficult to control)
- Feeling of having few reasons for living
- Demographic factors (male sex, indigenous or white ethnicity, middle to older age)
- Unmet acute or persistent physical health and behavioral health needs, including chronic pain, disability, substance use, and mood disorders
- Access to lethal means and gun ownership
- Social isolation and low belongingness
- Feeling hopeless about the future
- Unstable mood or sleeping patterns, including insomnia and nightmares
- Hospitalization or incarceration
- New or ongoing financial or employment problems

Suicide risk factors at the level of the relationship include:

- End of a relationship or marriage, including by death or divorce
- Relational problems, including abuse, and dissatisfaction.
- Unstable or conflictual relationships

Suicide risk factors at the level of the community include:

- Lack of access to appropriate and affirmative health and behavioral health care
- Disconnection from culture and cultural practices

Suicide risk factors at the level of the society include:

- Cultural beliefs or institutions that promote social isolation
- Sensationalistic media coverage, especially for youth
- Mental health stigma and discrimination

Protective Factors

Factors that lessen or protect against risk at the level of the person include:¹⁷⁴

- Life skills for coping, especially during stressful events and life changes (including problem solving skills and coping skills, ability to adapt to change)

- Personal or religious beliefs that prohibit or discourage suicide
- High self-esteem and sense of worth
- Strong quality of life with a purpose for living
- High sense of belongingness

Factors that lessen or protect against risk at the level of the relationship include:

- Connectedness to family or family of choice
- Genuine support from family or family of choice
- Relationships that affirm sexual identity and gender identity

Factors that lessen or protect against risk at the level of the community include:

- Access to appropriate and affirmative health and behavioral health care
- Connectedness to neighborhood, community, or social group
- Community members who check-in with one another
- Social institutions that promote healthy and active lifestyles

Factors that lessen or protect against risk at the level of the society include:

- Cultural or religious beliefs that prohibit or discourage suicide and value purposeful living
- Religious affiliation or spiritual community membership

Warning Signs

The following behaviors could indicate or signal suicide risk:¹⁷⁵

- Communicating wish to die or plans to attempt suicide
- Expressing the experience of suicidal thoughts that are intense, pervasive, or difficult to control
- Looking for a way to kill oneself, such as searching online or obtaining a gun
- Giving away possessions, or drafting notes indicating suicidal intent or desire
- Communicating feeling hopeless or having no reason to live or persistent hopelessness
- Communicating feelings of guilt, shame, or self-blame
- Communicating feelings of being trapped or in unbearable pain
- Communicating being a burden to others
- Increasing the use of alcohol or drugs
- Acting anxious or agitated; behaving recklessly or engaging in risky activities
- Insomnia, nightmares, and irregular sleeping
- Withdrawing or feeling isolated
- Communicating or exhibiting anxiety, panic or agitation
- Appearing sad or depressed or exhibiting changes in mood
- Showing rage or uncontrolled anger or communicating seeking revenge

Vulnerable Groups

Members of some groups and occupations may be more vulnerable to suicide than others. Despite this increased vulnerability, most people in the groups described below will not die by suicide or engage in suicidal behavior. Regardless of group membership, it is important to keep in mind suicide most often occurs among people with unmet behavioral health needs and is a symptom of depression.¹⁷⁶ This is not an exhaustive list and is intended to demonstrate differences and trends among groups and to highlight suicide prevention resources. Communities must utilize the Public Health Model to describe the problem of suicidal behavior and identify vulnerable community members, risk and protective factors, and effective interventions.

People in Middle and Older Age

Suicide rates among people in middle age – 35 to 64 years of age – are increasing.¹⁷⁷ Between 1999 and 2010, suicide rates among people in middle age have increased nearly 30 percent, especially among people aged 50 to 59.¹⁷⁸ In 2017, people of middle age represented 25.9 percent of the U.S. population but 35.1 percent of people who died by suicide.¹⁷⁹ Historically, older adults – or people over the age of 65 - have had the highest rates of suicide.¹⁸⁰ In 2017, this group represented 15.6 percent of the U.S. population but accounted for 18.2 percent of all suicides.¹⁸¹ The high suicide rates among older adults may be driven by factors such as use of highly lethal means; unmet health and behavioral health needs, especially late-life onset of depression; personality traits and coping mechanisms; life stressors, such as the loss of loved ones; social disconnection; and impairments in functioning and disability.¹⁸²

KEY RESOURCE: Preventing Suicide among Men in the Middle Years: Recommendations for Suicide Prevention Programs | Developed by the Suicide Prevention Resource Center: http://www.sprc.org/sites/default/files/resource-program/SPRC_MiMYReportFinal_0.pdf.

People Discharged from Hospital Settings

People seen in emergency departments for self-injury, regardless of their intent to die, are 30 times more likely to die by suicide than people who do not self-injure.¹⁸³ People discharged from psychiatric hospitalization are at especially high risk for future suicide and suicidal behavioral. Suicide risk increases during the first week of admission to a psychiatric hospital and during the first week after discharge.¹⁸⁴ For veterans, one study showed that suicide risk may be elevated during the first three months following discharge from a psychiatric hospital.¹⁸⁵ Common challenges that increase risk following discharge include missed follow-up appointments for outpatient care; a lack of resources or connection to such resources; unsupportive relationships or social networks, resulting in isolation and shame; and referrals that do not match individual needs.

KEY RESOURCE: Continuity of care for suicide prevention and research: Suicide attempts and suicide deaths subsequent to discharge from the emergency department or psychiatry inpatient unit | Developed by Knesper, D. J., American Association of Suicidology, & Suicide Prevention Resource Center: <http://www.sprc.org/sites/default/files/migrate/library/continuityofcare.pdf>.

Veterans

Veterans account for approximately 14 percent of all suicides in the U.S.¹⁸⁶ More than half of the veterans who die by suicide are 55 years of age or older, but the suicide rate among veterans between the ages of 18 and 34 has increased by 11 percent, rising from a rate of 40.4 deaths per 100,000 people in 2015 to 45 deaths per 100,000 people in 2016.¹⁸⁷ Data show that nearly 70 percent of veteran suicides are by firearm, compared to less than 50 percent of all non-veteran suicides.¹⁸⁸ This fact underscores the importance of considering the means by which vulnerable group members die by suicide in any suicide prevention strategy.¹⁸⁹ Veterans have unique risk and protective factors related to military service, in addition to factors previously mentioned.¹⁹⁰ Protective factors include a strong sense of belongingness to a unit and resilience to withstand adversity.¹⁹¹ On the other hand, transitioning out of military service may increase suicide risk.¹⁹² Stressful experiences during this transitional period include a loss of purpose and sense of identity, difficulties securing employment, conflicted relationships with family and friends, and other general challenges related to adapting to post-military life.¹⁹³

KEY RESOURCE: National Strategy for Preventing Veteran Suicide (2018-2028) | Developed by the U.S. Department of Veterans Affairs: https://www.mentalhealth.va.gov/suicide_prevention/docs/Office-of-Mental-Health-and-Suicide-Prevention-National-Strategy-for-Preventing-Veterans-Suicide.pdf.

Sexually and Gender Diverse People

“Sexually diverse” is an inclusive description referring to people who identify as lesbian, gay, bisexual, queer, and other sexual orientations.¹⁹⁴ While all members of this group may be vulnerable to suicide, research shows that people in lesbian, gay, and bisexual (LGB) communities may be at increased risk for suicidal behavior.¹⁹⁵ Community-based surveys estimate that 20 percent of LGB people have attempted suicide.¹⁹⁶ There are important differences among sexual orientation subgroups regarding suicide risk. Bisexual people report higher rates of suicidal ideation, followed by gays and lesbians.¹⁹⁷ Bisexual and questioning people – especially women – are at increased risk for depression, substance use, and suicidal behavior.¹⁹⁸ This may be due to feelings of invisibility and lack of community support, as bisexual people may experience rejection by both gay and straight communities, negative stereotypes, and feeling like they cannot be “out” about their sexuality.¹⁹⁹ Suicide risk also is elevated among transgender and gender non-conforming people.²⁰⁰ One study showed that 40 percent of transgender people attempted suicide at least once in their lifetime, with 92 percent

of those making the attempt before the age of 25.²⁰¹ Studies indicate that as many as 50 percent of transgender and gender non-conforming youth have attempted suicide.²⁰² Rejection of sexual and gender diversity by family and caregivers may significantly increase risk for suicide among LGBT youth, highlighting the need to include family-based interventions in suicide prevention efforts.²⁰³

KEY RESOURCE: Suicide risk and prevention for lesbian, gay, bisexual, and transgender youth | Suicide Prevention Resource Center: http://www.sprc.org/library/SPRC_LGBT_Youth.pdf.

Youth of Color

American Indian and Alaska Native youth and young adults have the highest rate of suicide compared to any other cultural or ethnic group in the United States.²⁰⁴ Suicide is the second leading cause of death for American Indian and Alaska Native children and adults ages 10 to 34.²⁰⁵ A recent study found African American children ages five to 12 – both boys and girls – are dying by suicide at twice the rate compared to white children.²⁰⁶ This finding highlights the need for continuous evaluation using the Public Health Model, as new at-risk groups emerge. Youth attempt suicide at greater rates compared to members of other age groups.²⁰⁷ Racial and ethnic differences also are found among suicidal behavior.²⁰⁸ Latina adolescents, in particular, report the highest rates of suicidal behavior compared to any other youth group.²⁰⁹ As many as one in seven Latina youth attempt suicide, a rate greater than any other youth group of the same age.²¹⁰

KEY RESOURCE: To Live To See the Great Day That Dawns: Preventing Suicide by American Indian and Alaska Native Youth and Young Adults | Developed by the U.S. Department of Health and Human Services: <https://store.samhsa.gov/system/files/sma10-4480.pdf>.

Rural Community Residents

People living in rural communities are at greater risk for suicide than those in more urban or densely populated communities.²¹¹ Rural communities may be more likely to feature characteristics with risk factors for suicide, such as gun ownership, social isolation, and difficulty accessing health and behavioral health care and social services.²¹² Even if services are available in rural communities, there are additional challenges that can affect the quality and timeliness of access.²¹³ These challenges include:

- A shortage of health care practitioners and providers to conduct preventative assessments and offer referrals and warm handoff to needed services, especially services that specifically address suicide risk
- Limited numbers of qualified, culturally competent providers and staff to deliver services
- Transportation, particularly in areas where people must travel long distances to seek services

- Insurance coverage that is accepted by the practitioner or provider
- Language barriers that prevent people from communicating with service providers
- Privacy concerns, especially for residents seeking mental health services in small communities²¹⁴

KEY RESOURCE: Understanding the Impact of Suicide in Rural America | National Advisory Committee on Rural Health and Human Services, Department of Health and Human Services: <https://www.hrsa.gov/sites/default/files/hrsa/advisory-committees/rural/publications/2017-impact-of-suicide.pdf>.

People Working in Certain Occupations

People in certain occupations are at increased risk for suicide.²¹⁵ Characteristics of occupations where risk might be elevated include jobs that are socially isolating; involve a high level of stress; are low paying or cause an increasing educational debt-to-income ratio; expose employees to violence or traumatic events; are fast-paced and require long hours; or are inconsistent, such as seasonal work.²¹⁶ The largest percentage – 20 percent in 2015 — of men who die by suicide are in construction and mining occupations.²¹⁷ Arts, design, entertainment, sports, and media occupations have the highest rates of suicide among both women and men. People in other occupations with increased risk include first responders, such as police, firefighters, and paramedics, as well as physicians, nurses, and veterinarians.²¹⁸

KEY RESOURCE: Comprehensive Blueprint for Workplace Suicide Prevention | National Action Alliance for Suicide Prevention: <https://theactionalliance.org/communities/workplace/blueprintforworkplacesuicideprevention>

People in Correctional Settings

People in correctional settings have higher rates of suicide compared to their counterparts in the community.²¹⁹ Correctional settings in California include prisons, jails, and juvenile detention facilities. Suicidal behavior may increase upon incarceration, but there is some evidence that people who become incarcerated may have experienced a history of suicidal behavior and other risk factors, such as unmet behavioral health needs, prior to becoming incarcerated.²²⁰ Elevated suicide risk also is found among people who work in correctional settings. One study found that correctional officers have a 39 percent higher chance of suicide compared to the average for other occupations.²²¹ This elevated risk for suicide may be due to work stress and its impact on family life, leading to separation and divorce.²²²

KEY RESOURCE: Suicide Prevention Resources for Adult Corrections | Developed by the Suicide Prevention Resource Center: <https://www.sprc.org/sites/default/files/resource-program/AdultCorrectionsResourceSheet.pdf>.

Best Practice in Suicide Prevention

The Institute of Medicine organizes suicide prevention activities along a continuum, ranging from universal to selective to indicated.²²³ Universal prevention efforts focus on the entire population and seek to deter suicidal behaviors by creating safe environments, increasing connectedness, building skills, and promoting mental health.²²⁴ Selective prevention efforts target people within vulnerable groups who have been identified as at greater risk for suicidal behaviors.²²⁵ Indicated prevention efforts focus on serving people engaged in suicidal behavior and providing timely intervention to prevent future suicidal behavior.²²⁶ Best practices reach across the social ecology, intersecting at person, relationship, neighborhood, and societal levels.²²⁷ Certain suicide prevention activities with strong evidence of effectiveness have demonstrated significant return on investment. These include training for health professionals; early identification of behavioral health needs, particularly depression; and creating barriers to prevent people from accessing methods to die by suicide.²²⁸

Best practices can only lead to successful outcomes if strong infrastructure is in place. For purposes of this plan, infrastructure refers to visible, multilevel leadership and networked partnerships; effective management of resources; and use of data for monitoring and improvement.²²⁹ Suicide prevention, as a public health challenge, is not unique in requiring infrastructure to support the delivery of best practices. An analysis of California's anti-tobacco initiative, for example, found that creating anti-smoking infrastructure was identified as the biggest challenge to the success of the effort.²³⁰ Many of the best practices described below are being delivered in select settings or communities throughout California.

Universal Prevention Strategies

Universal suicide prevention strategies are broad and are intended to reduce risk in the general population. Best practices in this category focus on the safety and health of the community through reducing access to lethal means, connecting people to social networks, building resiliency, safe reporting by the media following a suicide death, and increasing access to care. Research demonstrating the effectiveness of universal prevention strategies is scarce, limiting both knowledge about such strategies and investment in their development. The section below highlights best practices in universal suicide prevention.

Lethal Means Restriction

Lethal means restriction – or reducing someone's access to the lethal methods by which to die by suicide – is one of the best empirically supported methods of reducing suicide.²³¹ The effectiveness of reducing access to lethal means has been demonstrated in multiple countries and across a wide range of interventions.²³² The United Kingdom saw a reduction in suicides following replacement of coal gas – which contains carbon monoxide – with natural gas.²³³ After Israel adopted a policy requiring soldiers to lock their weapons in storage when on leave, suicide deaths were reduced by 40 percent.²³⁴ A ban on certain chemicals in Sri Lanka was

associated with a reduction in suicides involving pesticides in that country.²³⁵ Suicide deaths by carbon monoxide dramatically decreased following the implementation of strict controls on motor vehicle exhaust gas emissions in the U.S.²³⁶

Policies that limited the number of prescriptions written for certain medications, along with their pack size, resulted in decreased suicides involving those medications in several countries.²³⁷ Conversely, the potential consequences of removing safety measures also has been documented. The removal of safety barriers from a central city bridge in Australia, for example, led to an immediate increase in the numbers and rate of suicide at the bridge.²³⁸ Suicide deaths were reduced to zero at sites where barriers were removed and then reinstalled, as was the case in New Zealand.²³⁹ The effects of barrier installations are significant and immediate, and there is no evidence showing that their addition increases suicides at other locations or by other methods.²⁴⁰ In California, Caltrans is required to consider suicide risk in the design or redesign of bridges, and there are federal funds accessible for construction of suicide deterrent systems. However, there are no standards to guide prevention and policy at other sites.²⁴¹

The most effective methods of lethal means restriction are physical deterrents, which include carbon monoxide emission controls in vehicles; locking screen doors, windows, and drawers; suicide deterrent systems on railways and bridges; firearm safety mechanisms, such as gun locks and safes; and overdose prevention, such as the use of naloxone or blister packaging of medications.²⁴² Other effective methods include signage and connection to crisis services and means restriction counseling. Studies show that these methods can and should be combined with physical deterrents, where applicable.²⁴³

Focus on Common Lethal Means As demonstrated above, policies restricting the availability and accessibility of the means by which people die by suicide has the potential to significantly reduce suicide rates by those means. In California – and nationally – where suicide most commonly occurs when firearms are used, access and availability of firearms increases risk for unnatural death, including suicide.²⁴⁴ Firearms that are loaded or unlocked are tied to increased risk for intentional and unintentional death.²⁴⁵ Policies that reinforce gun safety and safe storage practices have been found to reduce risk for injury and death. For example, state bans on the sale of handguns that do not adhere to safety standards – sometimes referred to as “junk guns” – have demonstrated population-level effects on reducing suicide rates.²⁴⁶ Some states have expanded temporary transfer laws to include a temporary transfer a firearm from a person at risk to another person if such transfer is necessary to prevent imminent death or great bodily harm.²⁴⁷ Finally, research has shown an association between risk-based gun removal laws and a reduction in suicides by firearm.²⁴⁸ The Gun Violence Restraining Order is an example of a risk-based gun removal law in California.²⁴⁹ It is a court order that prohibits certain people – those experiencing suicidal or homicidal thoughts or behaviors, for example – from owning or buying guns and ammunition.²⁵⁰

In addition to policy changes to support means safety, programs to collaborate with gun shop and shooting range owners to prevent suicide among gun owners and their family members show promise. The Gun Shop Program, for example, was developed in New Hampshire after three people died by suicide by a firearm purchased at the same gun shop. Materials designed for and by gun shop owners were distributed to local shops and included information for identifying and interacting with a customer who may be at risk for suicide. Modeled after effective strategies in New Hampshire, the former Superior California Suicide Prevention Network developed best practice guidance for how to engage with community members on firearm suicide prevention messaging and approaches, such as increasing awareness of warning signs and increasing seeking help by people at risk.²⁵¹ Recognizing shared goals, the American Foundation for Suicide Prevention and the National Shooting Sports Foundation are collaborating to expand awareness of firearm safety measures to prevent suicide.²⁵² In Washington state, the National Rifle Association and the Second Amendment Foundation supported legislation to increase suicide prevention training and messaging for firearm professionals.

California Community Highlight: The Golden Gate Bridge's Suicide Deterrent System

California is home to several bridge and rail sites where people die by suicide in large numbers every year. The most well-known among these is the Golden Gate Bridge in San Francisco.

An average of 30 people die by suicide each year at the bridge site. Since the bridge opened in 1937, more than 1,700 people have lost their lives. Most people who die by suicide at the bridge are male, white, under 40 years of age, and live in the Bay Area. Fewer than 35 people have survived their attempt.

In addition to the roughly 30 known suicides in 2017, 235 people were saved from falling by a variety of public and private agencies and citizens, including the Golden Gate Bridge Patrol, California Highway Patrol, iron workers on the bridge, tow truck operators, Bridgewatch Angels volunteers, and many others.

Nets made of marine-grade woven steel, supported by scaffolding, are being installed to prevent death and deter people from considering the bridge a means of dying by suicide. The barrier will cost an estimated \$211 million in federal, state, and local funding.

Gaining approval to install the bridge barrier was not easy and took years, even requiring a change to federal transportation laws to allow for funding of suicide prevention projects. Many opponents of the bridge barrier cited aesthetic concerns. The barrier is expected to be fully installed by early 2021.

For more information, please visit <http://www.bridgerail.net/>.

Overdose is the most common method of suicide attempt.²⁵³ In addition to policies that restrict prescriptions and the amount of medications, other policies that increase the use of harm-reduction interventions can prevent overdose by certain drugs. For example, medication-assisted treatment, specifically naloxone, may reduce suicide by opioid overdose. Naloxone is a medication that works almost immediately to reverse opiate overdose. It has few known adverse effects, no potential for abuse, and can be rapidly administered through intramuscular injection or nasal spray. While most professional first responders and emergency departments are equipped with naloxone, emergency service providers may not arrive in time to revive overdose victims. In recent years, California has made naloxone more accessible through a statewide standing order allowing the administration of naloxone by family members and friends in a position to intervene during an opioid-related overdose.²⁵⁴

Assessing Access to Lethal Means Assessing access to lethal means and providing counseling to restrict such access are two best practices shown by evidence to reduce suicidal behavior.²⁵⁵ One study found that families of high-risk youth were significantly more likely to remove or secure lethal means in the home when counseled in the emergency department following suicidal behavior by a child.²⁵⁶ Despite such evidence, people identified as having suicidal ideation, or those who have been discharged from health care settings after attempting suicide, are not counseled routinely on means safety.²⁵⁷ Counseling on Access to Lethal Means (CALM) is a free resource available to identify people who could benefit from lethal means counseling, ask about their access to lethal methods, and work with them—and their families—to reduce access.²⁵⁸ Health care practitioners and providers are well-positioned to assess for access to lethal means when such a step is relevant to health care, but many feel uncomfortable doing so. In one study, community-based mental health providers were more likely to assess for and reduce access to lethal means collaboratively with people at risk and their families after they received training in CALM.²⁵⁹

Connectedness

Connectedness is the degree to which a person or group is socially close, interrelated, or shares resources with others.²⁶⁰ Connectedness can protect a person who is facing adversity. Peer programs in the military, for example, have been shown to effectively reduce risk for suicide when social networks are created between military members and their peers.²⁶¹ Although communities are not necessarily bound by neighborhoods, schools, or other institutions, these structured environments can be catalysts for reducing suicide risk among a broad population. School connectedness has consistently been shown to play a critical role in protecting adolescents against many negative outcomes, including suicidal behaviors.²⁶² Groups that promote connectedness, such as the school-based Gay-Straight Alliance, show promise in reducing suicidal ideation and attempt among youth.²⁶³ Family connectedness can buffer against suicide risk. Family acceptance of sexual and gender identity among youth has been demonstrated to protect against suicide risk, and can be modified using evidence-based approaches, such as the Family Acceptance Project's Family Intervention Approach.²⁶⁴

Risk for suicide is reduced when people have trust in social networks and are engaged in community.²⁶⁵ Research shows that there is a relationship between connectedness and safety, namely that people are more likely to socially engage in environments that are safe, affirmative, supportive, and free of violence and discrimination.²⁶⁶ Suicidal behavior may share risk and protective factors with other forms of violence, such as domestic violence and children and elder maltreatment.²⁶⁷ Shared risk factors include lack of social support, economic stress, and substance use.²⁶⁸ Shared protective factors include the coordination of community resources and services, connectedness, and family support.²⁶⁹ Prevention resources to create training, programs, and partnerships can be used collectively to respond to multiple forms of violence, including suicide.²⁷⁰ Addressing multiple forms of violence is a prudent approach, especially since different forms of violence overlap and intersect.²⁷¹

Resilience and Skills-training

Resilience is the ability to withstand, adapt to, and recover from adversity, threat, and stress. Resilience is associated with coping, or people's individualized ability to manage both everyday stressors as well as more extreme stressors in their lives. Communities – including neighborhoods, schools, and organizations – can build resilience by strengthening cultural values and cultural identity; by reinstituting collective history, language, spirituality, and healing practices; and through collective action.²⁷² Culture in this context can refer to racial/ethnic; vocational, such as first responder and culinary; and special population, such as military culture.

Effective life skill interventions include techniques that promote critical thinking, conflict resolution, stress management, and coping and that help people safely manage challenges such as economic stress, divorce, physical illness, and aging. Best practice approaches to building universal life skills have been developed for school-aged children and youth. The Good Behavior Game, for example, is an early education classroom management technique that shows promise in reducing suicidal behavior for decades following program delivery.²⁷³ Life skills programs tailored to specific cultural norms and values also are supported by evidence of their effectiveness. One, the American Indian Life Skills Development curriculum, shows promise in reducing depression and suicidal behavior among Native youth.²⁷⁴

Responsible Media Reporting

Exposure to suicidal behavior by one person may facilitate the occurrence of subsequent, similar behaviors by others, especially among adolescents.²⁷⁵ Due to exposure, multiple suicides may occur within a particular time period or location, a pattern known as a suicide cluster.²⁷⁶ Suicide clusters are rare and happen almost exclusively among youth.²⁷⁷ The media may inadvertently increase suicide risk when reporting the details of a suicide.²⁷⁸ For example, extensive media coverage of suicide – in amount, duration, and prominence – is associated with increases in suicide rates.²⁷⁹ Harmful media practices, such as reporting details about the method used, also may increase risk for suicidal behavior in others, especially young people.²⁸⁰ Further, suicidal behavior using a particular method – even an uncommon method – may increase if that method is identified and described in media reports.²⁸¹

Best practice for responsible reporting of suicide include communicating messages demonstrating that suicide is preventable, printing or airing stories of hope and resilience, providing links to helping resources, and refraining from airing or publishing reports that sensationalize suicide. Local media can partner in effective suicide prevention by disseminating the message that suicide is preventable through fictional story lines, real-life reporting, billboards, and public service announcements.²⁸² Positive storylines about mental health and suicide can prompt media consumers to take direct action to seek or provide help.²⁸³ Such storylines also empower people to have open conversations with friends and family.²⁸⁴

California Community Highlight: Response Following Suicide Cluster

Between May 2009 and March 2015, nine people who were either incoming or current high school students or alumni of a single Santa Clara County school district died by suicide. The California Department of Public Health requested assistance from the Centers for Disease Control and Prevention and the Substance Abuse and Mental Health Services Administration to investigate the deaths and understand how youth suicide in Santa Clara County, its school districts, and cities could be prevented in the future.

Recommendations included:

1. Using multiple prevention approaches to address multiple risk factors
2. Ensuring access to evidence-based mental health care
3. Strengthening family relationships and family-based programs
4. Increasing students' connection to school and school-based programs
5. Identifying and supporting people at risk
6. Strengthening crisis Intervention
7. Delivering services to loss survivors in the event of a student suicide
8. Launching prevention efforts involving other forms of violence
9. Reducing access to lethal means for youth at-risk
10. Using safe messaging and reporting about suicide
11. Engaging in strategic planning for suicide prevention
12. Selecting and implementing evidence-based programs
13. Mandating continuous program evaluation

For more information, please visit <https://www.sccgov.org/sites/phd/hi/hd/epi-aid/Documents/epi-aid-report.pdf>.

Access to Health and Behavioral Health Care

Services that deliver appropriate, timely, and accessible health and behavioral health care have the potential to prevent suicide. Best practices include administrative policies, such as full coverage of behavioral health needs in insurance policies and managed care, as well as policies

that address provider shortages, especially in rural and underserved communities.²⁸⁵ Policies to address provider shortages include the use of financial incentives and expansion of telehealth approaches that connect providers and clients through phone, video, and internet-based technologies.²⁸⁶ Mobile and telehealth approaches may increase access to health care, especially in physically isolated communities.²⁸⁷ Research on telehealth approaches to suicide care is limited but promising.²⁸⁸

Clear messaging to create easy pathways to available services also shows promise for suicide prevention. Messaging to support people to seek help includes teaching early recognition of behavioral health needs and reducing stigma associated with seeking help by normalizing the behavior among peers. Peer norm programs seek to normalize protective factors – including help-seeking and reaching out and talking to trusted people – and also promote peer connectedness.²⁸⁹ By leveraging the leadership qualities and social influence of peers, these approaches can be used to shift group-level beliefs and promote positive social and behavioral change.²⁹⁰ This approach has been especially successful in school settings but also has shown promise in the workplace and other settings.²⁹¹

Selective Prevention Strategies

Selective prevention strategies are those focused on detection of risk and screening select subgroups that may develop risk for suicidal behaviors. Best practices in this category are effective strategies used to identify risk and intervene early, and to connect people to services. Best practices in selective suicide prevention are highlighted below.

Collaborative Care

Collaborative care is an integrated care model that has been tested in over 80 randomized control trials. While it has not specifically been shown to reduce suicide, studies have confirmed the benefits of collaborative care for people with risk factors for suicide, namely depression and anxiety.²⁹² Under this model, traditional primary care is integrated with a team comprised of a care coordinator and a specialty behavioral health provider.²⁹³ This team collaborates to create a holistic plan for the person based on best practices, client-directed goals, and the monitoring of those goals, making adjustments as needed when progress is stalled. Two landmark studies demonstrate reduced suicidal ideation using collaborative interventions for older adults experiencing depression. The Prevention of Suicide in Primary Care Elderly: Collaborative Trial reduced suicidal ideation and depression among older adults through a collaborative approach between a person, a primary care physician, and a health specialist, such as a nurse, social worker, or mental health provider.²⁹⁴ Second, the Improving Mood—Promoting Access to Collaborative Treatment approach involves collaboratively developing a care plan – with input from the person, primary care provider, care manager, and consulting psychiatrist – to reduce depression and suicidal ideation in older adults. Evaluation of this model demonstrated

significant decreases in depression and suicidal ideation, in addition to improved functional and quality of life outcomes.²⁹⁵

Depression Screening and Management by Physicians

The majority of people who die by suicide had contact with their primary care physician in the year prior to death, while almost half had contact in the month preceding death.²⁹⁶ Despite such contact, suicide risk is under-recognized and under-served in these critical primary care settings.²⁹⁷ Nearly 70 percent of people experiencing depression who see a primary care physician will report physical complaints, such as physical pain or sleep disturbances.²⁹⁸ Training for primary care physicians on identification of suicide risk and treatment of depression and other risks, such as substance use, shows promise in preventing suicide, especially when delivered in collaborative care models.²⁹⁹

Gatekeeper Training

Gatekeeper training is designed to train teachers, families, coaches, military commanders, supervisors, clergy, emergency responders, urgent care providers, and others in the community to identify people who may be at risk of suicide and to respond effectively, including facilitating connection to services.³⁰⁰ Gatekeeper training focuses on increasing a person's ability to recognize warning signs of suicide and provide referral.³⁰¹ Some trainings include information on delivering brief interventions to support people at risk for suicide, such as reducing a person's access to lethal means.³⁰² Gatekeeper training may be implemented in a variety of settings to identify and support people at risk.³⁰³ Gatekeeper trainings have been shown to increase knowledge of risk factors and warning signs and increase confidence among people responding to someone expressing a desire to die.³⁰⁴

California Community Highlight: Available Gatekeeper Trainings

Below are options for suicide prevention awareness and support trainings for gatekeepers. This list is non-exhaustive and is intended to give the reader a starting point to explore available trainings.

Question, Persuade, Refer (QPR) | <https://qprinstitute.com/>.

Trainings by Living Works | <https://www.livingworks.net/>.

Trainings specific to school-settings available through the American Foundation for Suicide Prevention | <https://afsp.org/our-work/education/more-than-sad/> and <https://afsp.org/our-work/education/signs-matter-early-detection/>.

Crisis Response

Crisis response can include a variety of crisis services, ranging from warm lines and crisis lines to crisis stabilization support and short-term crisis residential care.³⁰⁵ Best practice approaches for systematic crisis response include centralized call centers that use real-time coordination across

systems, coordinated mobile crisis outreach and support, and crisis residential and stabilization services.³⁰⁶ The delivery of coordinated crisis services also has been shown to reduce redundancies and costs associated with connecting people with an appropriate level of care to prevent suicidal behavior.³⁰⁷

Under effective models, suicide prevention hotline, text, and chat services provide 24-hour support to conduct suicide assessment and intervention, provide referrals to appropriate services, help people develop safety plans, and connect people with mobile crisis or emergency resources.³⁰⁸ The hotlines generally prevent suicide in two ways: They ensure the immediate safety of at-risk callers, and they link those who may be at risk of suicide with appropriate and available resources.³⁰⁹ Effective training and standards for practice are critical. A study of crisis line staff who received Applied Suicide Intervention Skills Training showed improved outcomes for callers, including reduced depression, a reduced sense of being overwhelmed, lower suicide risk, and increased hopefulness.³¹⁰

California Community Highlight: Caltrans District 7 and Didi Hirsch Collaboration

Local transportation leaders are partnering with suicide prevention centers to create safe environments with physical deterrents and crisis services messaging and response. Caltrans District 7, which covers Los Angeles and Ventura counties, in partnership with Didi Hirsch Mental Health Services and regional first responders, are working to prevent suicide by identifying community sites used for suicidal behavior, constructing barriers, when feasible, and installing suicide hotline signage and cameras, where appropriate. The effort is supported by a committed network of partners, including first responders, facility and equipment owners, suicide prevention and crisis services, and local authorities. Coordination continues once a site is identified and fortified. For example, trained camera monitors identify a person at risk and alert first responders and crisis services.

For more information, please visit <https://didihirsch.org/>.

Indicated Prevention Strategies

Indicated prevention strategies focus on people engaged in suicidal behavior and people bereaved by the loss of a loved one to suicide. Best practices in this category focus on providing care that specifically targets suicidal behavior and following-up with people who have been discharged from health care settings after being served for suicidal behavior. Indicated prevention best practices also deliver coordinated, timely, and respectful services to suicide loss survivors.

Suicide Risk Assessment and Management

Best practice for screening and risk assessment in health and behavioral health care settings includes provider knowledge of risk and protective factors and warning signs, procedures for categorizing risk and making clinical decisions based on risk, evidence-based assessments and safety planning, documentation of risk level and action taken, and caring referral procedures.³¹¹ Standardization makes the entire process of identifying risk and connecting people to services transparent and collaborative for the provider and person at risk.³¹² Critical to this collaborative process is obtaining informed consent and the use of a standardized decision-making process to routinize risk designations based on suicide attempt history, the severity of current suicidal symptoms, and the integration of risk factors.³¹³ Standardizing risk assessment and management has the potential to reduce clinical or legal concerns about errors in judgement that might overestimate or underestimate risk.³¹⁴ Suicide risk assessments help identify acute, modifiable, and treatable risk factors and help providers recognize when people need more structured methods for managing daily living.³¹⁵

The Columbia-Suicide Severity Rating Scale is a common screening tool that uses a series of questions in plain language to help users identify whether a person is at risk for suicide, and also assess the severity and immediacy of the risk, and identify possible support.³¹⁶ The tool is suitable for all ages and special populations and is available in over 100 country-specific languages.³¹⁷ In health care settings, the Patient Health Questionnaire (PHQ9) is an assessment that asks nine questions about depressive symptoms experienced in the prior two weeks, with one question devoted to thoughts of dying or being better off dead. The PHQ9A is the PHQ9 modified for adolescents ages 11 to 17.³¹⁸ Finally, the Ask Suicide-Screening Questions is a tool used to identify a youth at risk in medical settings and takes less than one minute to complete.³¹⁹ Positive screens obtained through use of this tool prompt providers to conduct additional, in-depth assessments.³²⁰

Safety planning is a brief intervention that incorporate best practices in means restriction, problem-solving, social support, and emergency resources.³²¹ Safety planning is not a “no-harm contract” or “contract for safety” that requires people at risk to promise a provider the person will not engage in suicidal behavior; research shows such “contracts” are not effective and

actually can increase risk.³²² The Stanley Brown Safety Plan and Crisis Response Planning tools are evidence-based and commonly used in many settings. The Stanley Brown Safety Plan includes methods for keeping homes safe, recognizing warning signs of a suicidal crisis, identifying ways to cope with suicidal thoughts, and identifying friends, family, and mental health and emergency resources, such as the location of the nearest emergency department.³²³ Crisis Response Planning is a strategy used to develop written steps for a person at risk for suicide to take during times of crisis or when under stress. Using an index card, people list steps for identifying personal warning signs, along with coping strategies and social and professional support. Results of a randomized clinical trial show that crisis response planning reduced suicide attempts by 75 percent compared to using safety contracts, or contracts in which a person vows not to self-injure.³²⁴

Treatment Interventions

Effective care that targets suicide risk, specifically, is effective when it is structured and integrates problem-solving skills, collaborative assessment, service planning, and caring, consistent follow-up.³²⁵ Below are behavioral and pharmacological interventions shown to be efficacious in the treatment of suicidal behaviors:

- **Dialectical Behavioral Therapy** is a cognitive-behavioral treatment that combines therapy, skills training, and coaching and has been shown to be effective for treating suicidal behavior and non-suicidal self-injury at any age.³²⁶ Dialectical Behavioral Therapy has been adapted for adolescents in a shorter format – from 12 months to 16 weeks – and includes skill modules to improve parent-child communication, among other skills.³²⁷ In addition, nonclinical applications have been adapted for school-settings and teach students in grades 6-12 mindfulness, emotional regulation, and interpersonal skills.³²⁸
- **Cognitive Behavioral Therapy for Suicide Prevention** is a cognitive behavioral treatment for people who have attempted suicide within the last 90 days.³²⁹ The primary goals of this intervention are to reduce suicidal risk factors, enhance coping skills, and prevent future suicidal behavior.³³⁰ The therapy is designed to help people use more effective means of coping with stressors and problems that trigger suicidal crises.³³¹
- **Collaborative Assessment and Management of Suicidality** is a suicide-specific therapeutic framework that can be delivered with other treatments and across different settings, including community and inpatient settings.³³² A psychotherapeutic framework that “amplifies active collaboration” between a service provider and a person at risk, it assesses for and addresses factors that are increasing risk.³³³ The alliance between provider and client is intended to support the person at risk’s motivation to live.³³⁴
- **The Attempted Suicide Short Intervention Program (ASSIP)** is a brief intervention specifically for attempt survivors.³³⁵ It emphasizes the therapeutic alliance between provider and survivor developed in an initial interview. Findings are promising. When

combined with clinical treatment, ASSIP was able to reduce suicidal behavior over a two-year period for people who recently attempted suicide.³³⁶ ASSIP also has been demonstrated to reduce health care costs.³³⁷

- **Pharmacological interventions** can reduce suicide risk by addressing mental health needs.³³⁸ Antidepressants, such as selective serotonin reuptake inhibitors, can alleviate depression and associated suicide risk.³³⁹ Lithium for the treatment of mood disorders and clozapine for the treatment of schizophrenia have been shown to reduce suicide among people receiving services for people with these needs.³⁴⁰

Innovations in this area are underway, and target highly treatable risk factors – such as insomnia - using low-risk interventions to prevent suicide.³⁴¹ Non-mental health interventions show promise for targeting risk. One example is services that address sleep disturbances, which may reduce risk and can be delivered through brief, targeted interventions.³⁴² Repetitive transcranial magnetic stimulation (rTMS) also shows promise in addressing suicidal ideation. This approach uses a magnet to target and stimulate specific areas of the brain and is typically used to treat depression and anxiety. In one study, 40 percent of people served with bilateral rTMS therapy reported no longer experiencing thoughts of suicide.³⁴³ In addition, ketamine is a pharmaceutical drug recently approved for therapeutic use to rapidly reduce depressive symptoms and suicidal ideation.³⁴⁴ Acute suicide risk is almost immediately reduced, and beneficial effects can extend up to 10 days.³⁴⁵

Emergency Department Interventions

Emergency departments play a key role in suicide prevention efforts.³⁴⁶ Statistics show that 20 percent of people who die by suicide visited an emergency department within a month of death, and 60 percent of survivors of suicide attempt sought medical care for their injuries in emergency departments. National data suggest that interventions in the emergency department may decrease suicide deaths by 20 percent.³⁴⁷ The Emergency Department Safety Assessment and Follow-Up Evaluation study evaluated an emergency department intervention that combined universal screening for suicide risk; secondary assessment by a physician; resources at discharge, including a safety plan; and follow-up telephone calls over a year-long period. The study found significant decreases in suicidal behavior among people who received the intervention.³⁴⁸

The effectiveness of delivering follow-up care – also referred to as caring contacts – to people discharged from hospital settings after suicidal behavior is backed by strong evidence.³⁴⁹ One of the most empirically successful approaches to suicide prevention was the “caring letters study,” in which contact after discharge significantly reduced suicide among people who were hospitalized for depression or suicide risk.³⁵⁰ People who participated in the study were contacted using low-cost methods, such as postcards and short caring notes, at least four times a year for five years.³⁵¹ Suicide rates were compared with those for people who received no contact following discharge during the same period.³⁵² People in the contact group had a lower

suicide rate in all five years of the study.³⁵³ Another study demonstrated significant return-on-investment for commercial insurance and managed care plans when people released from hospital or emergency departments for suicidal behavior received follow-up phone calls.³⁵⁴ Follow-up calls from crisis line providers are cost-effective, and have been demonstrated to reduce future suicidal behavior for people discharged from health care settings.³⁵⁵

California Community Highlight: WellSpace Health

California communities are linking suicide prevention centers with health care systems to deliver best practices. One example is WellSpace Health in Sacramento. WellSpace Health delivers integrated health and behavioral health care and operates the Suicide Prevention Crisis Line serving Northern California counties. One program, the Primary Care Follow Up Suicide Prevention program integrates screening for suicide risk in primary care and refers people to the 24-hour crisis lines through the electronic health record, and provides 30 days of follow-up, risk monitoring, emotional support, resource linkage, and safety planning. Another program, the Emergency Department Follow-Up program, connects with people at risk who are nearing discharge from hospital settings within 24 hours of discharge and delivers follow-up services, include emotional support, risk assessment, safety planning, and monitoring.

For more information, please visit <https://www.wellspacehealth.org/services/behavioral-health-prevention/suicide-prevention>.

Postvention

Postvention efforts are organized prevention activities directed toward suicide loss survivors, or people who have lost a loved one to suicide. These survivors may include family, friends, clinicians, physicians, coworkers, and crisis line volunteers. Loss survivors may encounter stigma associated with suicide, a reaction that may not accompany other manners of death and can act as a profound barrier to overcoming grief.³⁵⁶ Activities that may carry benefits for loss survivors include services to address grief and distress associated with suicide loss, services that specifically mitigate negative effects of exposure to suicide, and services that prevent suicide by people at risk following exposure to suicide.³⁵⁷ Face-to face bereavement support groups are the most studied intervention for loss survivors, while bereavement services that take a family-oriented approach show promise.³⁵⁸ With this model, family members can explore together their individual responses following a suicide and assess the family's collective response.³⁵⁹ Family members may become more engaged in the healing process because the family support system is also being served and potential miscommunication or dysfunction is reduced.³⁶⁰

References

- ¹ World Health Organization. (2014). *Preventing suicide: A global imperative*. Luxembourg: Author.
- ² Joiner, T. E., Jr. (2005). *Why people die by suicide*. Cambridge, MA: Harvard University Press.
- ³ Ibid.
- ⁴ Owens, D., Horrocks, J., & House, A. (2002). Fatal and non-fatal repetition of self-harm: Systematic review. *British Journal of Psychiatry*, 181, 193-199.
- ⁵ Talseth, A. G., Jacobsson, L. & Norberg, A. (2001). The meaning of suicidal psychiatric inpatients' experiences of being treated by physicians. *Journal of Advanced Nursing*, 34(1), 96-106.
- ⁶ World Health Organization (2014). *Preventing suicide: A global imperative*. Luxembourg: Author.
- ⁷ Hedegaard, H., Curtin, S. C., & Warner, M. Suicide mortality in the United States, 1999-2017. NCHS data brief, 330, 1-8. Hyattsville, MD: National Center for Health Statistics. 2011. Retrieved November 29, 2018 from <https://www.cdc.gov/nchs/data/databriefs/db330-h.pdf>.
- ⁸ Crosby, A. E., Han, B., Ortega, L. A. G., Parks, S. E., Gfoerer, J. (2011). Suicidal thoughts and behaviors among adults aged ≥ 18 years - United States, 2008-2009. *MMWR Surveillance Summaries*, 60 (SS-13), 1-22. Retrieved November 29, 2018 from www.cdc.gov/mmwr/preview/mmwrhtml/ss6013a1.htm?s_cid=ss6013a1_e.
- ⁹ Goldsmith, S. K., Pellmar, T. C., Kleinman, A. M., & Bunney, W. E. (Eds.). (2002). *Reducing suicide: A national imperative*. Washington, DC, US: National Academies Press.
- ¹⁰ Substance Abuse and Mental Health Services Administration. (2017). *Key substance use and mental health indicators in the United States: Results from the 2016 National Survey on Drug Use and Health* (HHS Publication No. SMA 17-5044, NSDUH Series H-52). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from <https://www.samhsa.gov/data/>.
- ¹¹ Shepard, D. S., Gurewich, D., Lwin, A. K., Reed, G. A., & Silverman, M. M. (2016). Suicide and suicidal attempts in the United States: Costs and policy implications. *Suicide & Life-Threatening Behavior*, 46(3), 352–362.
- ¹² Centers for Disease Control and Prevention. (2019). *Data & Statistics (WISQARS™): Cost of injury reports*. Retrieved June 18, 2019 from <https://wisqars.cdc.gov:8443/costT/>.
- ¹³ Bernert, R. A. (2018). *Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California*. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved December 14, 2018 from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ¹⁴ World Health Organization. (2012). *Public health action for the prevention of suicide: A framework*. Geneva, Switzerland: Author.
- ¹⁵ Ibid.

-
- ¹⁶ Office of Disease Prevention and Health Promotion. (n.d.). *HealthyPeople2020*. Retrieved April 29, 2019 from <https://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-of-health>.
- ¹⁷ World Health Organization. (2014). *Preventing suicide: A global imperative*. Luxembourg: Author.
- ¹⁸ World Health Organization. (2018). *National suicide prevention strategies: progress, examples and indicators*. (License: CC BY-NC-SA 3.0 IGO). Geneva: Author.
- ¹⁹ Carr, C. (2018). Presentation to the Mental Health Services Oversight and Accountability Commission in San Leandro, California on October 25, 2018.
- ²⁰ Ibid.
- ²¹ Ibid.
- ²² Ibid.
- ²³ Visit <https://resources.depaul.edu/abcd-institute/Pages/default.aspx> for more information on Asset-Based Community Development.
- ²⁴ Preti, A., Tondo, L., Sisti, D., Rocchi, M., & Girolamo, B. (2010). Correlates and antecedents of hospital admission for attempted suicide: a nationwide survey in Italy. *European Archives of Psychiatry and Clinical Neuroscience*, 260(3), 181–190.
- ²⁵ Chu, J. P., Goldblum, P., Floyd, R., & Bongar, B. (2010). The cultural theory and model of suicide. *Applied and Preventative Psychology*, 14 (1-4), 25-40.
<https://doi.org/10.1016/j.appsy.2011.11.001>.
- ²⁶ The Joint Commission (2018). *Requirement, rationale, reference report*. Issue 18, November 27, 2018. Retrieved from https://www.jointcommission.org/assets/1/18/R3_18_Suicide_prevention_HAP_BHC_11_27_18_FINAL.pdf.
- ²⁷ Little, V., Neufeld, J., & Cole, A. R. (2018). Integrating safety plans for suicidal patients into patient portals: Challenges and opportunities. *Psychiatric Services*, 69(6), 618-619.
<https://doi.org/10.1176/appi.ps.201700458>.
- ²⁸ Visit <https://www.211la.org/mayors-challenge> for more information on the Los Angeles Mayor's Challenge.
- ²⁹ California Department of Mental Health. (2012). *Transition plan: Transfer of the Department of Mental Health's Community Mental Health Programs to other state departments and organizations*. Retrieved January 2, 2019 from http://www.dsh.ca.gov/Publications/docs/Transition_Plan/DMHTransitionPlan.pdf.
- ³⁰ Department of Health Care Services. (2018). *Mental Health Services Act Expenditure Report – Governor's Budget: Fiscal Year 2018-19*. Retrieved January 2, 2019 from https://www.dhcs.ca.gov/formsandpubs/Documents/Legislative%20Reports/Mental%20Health/MHSA_Expenditure_Report-Jan2018.pdf.
- ³¹ Title 9 California Code of Regulations, Division 1, Chapter 14 MHSA, Section 3200.245. Prevention and Early Intervention Component.
- ³² Data compiled by MHSoAC staff using the MHSoAC Transparency Suite retrieved February 26, 2019 from <http://mhsoac.ca.gov/mhsoac-transparency-suite>.
- ³³ Visit <http://www.stanislausmhsa.com/pdf/public/annualupdates/mhsafy1819final.pdf> for more information on Stanislaus County's Suicide Prevention Innovation Project.

-
- ³⁴ Results cited in a document titled *Recommendation to the State Superintendent of Public Instruction* provided during the February 12, 2019 meeting of the California Department of Education's Student Mental Health Policy Workgroup.
- ³⁵ Chaptered by Secretary of State. Chapter 182, Statutes of 2017.
- ³⁶ Chaptered by Secretary of State. Chapter 527, Statutes of 2018.
- ³⁷ Chaptered by Secretary of State. Chapter 32, Statutes of 2018.
- ³⁸ Department of Mental Health Information Notice No. 08-25, Enclosure 1.
- ³⁹ Visit <https://calmhsa.org/> for more information about the California Mental Health services Authority.
- ⁴⁰ Clark, W., Collentine, A. M., Welch, S. N., & Brichler, S. (2013). *Best practices and initial outcomes of California's historic effort to reduce stigma of mental illness, prevent suicide, and improve student mental health. Presentation at the American Public Health Association.* Retrieved from <https://calmhsa.org/wp-content/uploads/2013/10/Final-APHA-Uploaded-Presentation.pdf>.
- ⁴¹ Visit <https://www.suicideispreventable.org/> for information about the Know the Signs Campaign.
- ⁴² Visit <http://www.directingchange.ca.org/> for information about the Direction Change Program.
- ⁴³ California Department of Mental Health. (2008). *California strategic plan on suicide prevention: Every Californian is part of the solution.* Retrieved January 4, 2019 from https://www.sprc.org/sites/default/files/California_CalSPSP_V92008.pdf.
- ⁴⁴ Department of Health Care Services. (2016). *Suicide Hotline Report.* Retrieved January 4, 2019 from https://www.dhcs.ca.gov/formsandpubs/Documents/Legislative%20Reports/2016Suicide_Hotline_Report.pdf.
- ⁴⁵ Ibid.
- ⁴⁶ Ibid.
- ⁴⁷ Department of Health Care Services. (2019). *Mental Health Services Act Expenditure Report – Governor's Budget, Fiscal Year 2019-20.* Retrieved March 15, 2019 from https://www.dhcs.ca.gov/formsandpubs/Documents/Legislative%20Reports/Mental%20Health/MHSA_Expenditure_Report-Jan2019.pdf.
- ⁴⁸ Visit <https://www.cdph.ca.gov/Programs/CHSI/Pages/End-of-Life-Option-Act-.aspx> for more information on California's End of Life Option Act.
- ⁴⁹ Crosby, A. E., Ortega, L., & Melanson, C. (2011). *Self-directed Violence Surveillance: Uniform Definitions and Recommended Data Elements*, Version 1.0. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control.
- ⁵⁰ Ibid.
- ⁵¹ Ibid.
- ⁵² Nock, M. K., Hwang, I., Sampson, N. A., & Kessler, R. C. (2010). Mental disorders, comorbidity and suicidal behavior: Results from the National Comorbidity Survey Replication. *Molecular Psychiatry*, 15(8), 868–876.
- ⁵³ Posner, K., Oquendo, M. A., Gould, M., Stanley, B., & Davies, M. (2007). Columbia Classification Algorithm of Suicide Assessment (C-CASA): classification of suicidal events in the

FDA's pediatric suicidal risk analysis of antidepressants. *The American Journal of Psychiatry*, 164(7), 1035-1043.

⁵⁴ Ibid.

⁵⁵ United States Food and Drug Administration, United States Department of Health and Human Services. (August 2012). *Guidance for Industry: Suicidality: Prospective Assessment of Occurrence in Clinical Trials, Draft Guidance. Revision 1*. Silver Spring, MD: Center for Drug Evaluation and Research.

⁵⁶ Jacobs, D. (2009). *Suicide Assessment Five-Step Evaluation and Triage for mental health professionals (SAFE-T)*. Retrieved from https://www.integration.samhsa.gov/images/res/SAFE_T.pdf.

⁵⁷ Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... Mann, J. J. (2011). The Columbia-Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *The American journal of psychiatry*, 168(12), 1266–1277. <https://doi.org/10.1176/appi.ajp.2011.10111704>.

⁵⁸ Bridge, J.A., Goldstein, T.R., & Brent, D.A. (2006). Adolescent suicide and suicidal behavior. *J Child Psychol Psychiatry*, 47(3–4), 372–394.

⁵⁹ Van Orden, K. A., Witte, T. K., Cukrowicz, K. C., Braithwaite, S. R., Selby, E. A., & Joiner, T. E. (2010). The interpersonal theory of suicide. *Psychological review*, 117(2), 575-600.

⁶⁰ Joiner, T. E., Jr. (2005). *Why people die by suicide*. Harvard University Press.

⁶¹ Baumeister, R., & Leary, M.R. (1995). The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychological Bulletin*, 117(3), 497-529.

⁶² Joiner, T. E., Jr. (2005). *Why people die by suicide*. Harvard University Press.

⁶³ Joiner, T. E., Hollar, D., Van Orden, K. A. (2006). On Buckeyes, Gators, Super Bowl Sunday, and the Miracle on Ice: "Pulling Together" is associated with lower suicide rates. *Journal of Social and Clinical Psychology*, 25, 180–196.

⁶⁴ Ibid.

⁶⁵ Ibid.

⁶⁶ Joiner, T. E., Jr. (2005). *Why people die by suicide*. Harvard University Press.

⁶⁷ Ibid.

⁶⁸ Ibid.

⁶⁹ Ibid.

⁷⁰ Joiner, T. E. (2010). Overcoming the fear of lethal injury: Evaluating suicidal behavior in the military through the lens of the interpersonal–psychological theory of suicide. *Clinical Psychology Review*, 30(3), 298-307. <https://doi:10.1016/j.cpr.2009.12.004>.

⁷¹ Yip, P.S., Caine, E., Yousuf, S., Chang, S., Wu, K.C., & Chen, Y. (2012). Means restriction for suicide prevention. *Lancet*, 379(9834), 2393-2399.

⁷² Sarchiapone, M., Mandelli, L., Iosue, M., Andrisano, C., & Roy, A. (2011). Controlling access to suicide means. *International journal of environmental research and public health*, 8(12), 4550–4562. doi:10.3390/ijerph8124550.

⁷³ Clarke, R., & Lester, D. (1989). *Suicide: closing the exits*. New York, NY: Springer Verlag.

⁷⁴ Hawton K. (2007). Restricting access to methods of suicide. *Crisis*, 28(S1), 4-9.

⁷⁵ Owens, D., Horrocks, J., & House, A. (2002). Fatal and non-fatal repetition of self-harm: Systematic review. *British Journal of Psychiatry*, 181, 193-199.

-
- ⁷⁶ Barber, C. W., & Miller, M. J. (2014). Reducing a suicidal person's access to lethal means of suicide: A research agenda. *American Journal of Preventative Medicine*, 47(3), S264-S272.
- ⁷⁷ Drexler, M. (n.d.) *Guns and suicide: The hidden toll*. Harvard Public Health: Magazine of the Harvard T.H. Chan School of Public Health. Retrieved November 8, 2018 from https://www.hsph.harvard.edu/magazine/magazine_article/guns-suicide/.
- ⁷⁸ Ibid.
- ⁷⁹ Ibid.
- ⁸⁰ Brown, G. K., Henriques, G. R., Sosdjan, D., & Beck, A. T. (2004). Suicide intent and accurate expectations of lethality: predictors of medical lethality of suicide attempts. *Journal of Consulting and Clinical Psychology*, 72(6), 1170-1174.
- ⁸¹ Bostwick, J. M., Pabbati, C., Geske, J. R., & McKean, A. J. (2016). Suicide attempt as a risk factor for completed suicide: Even more lethal than we knew. *The American Journal of Psychiatry*, 173, 1094–1100. <https://doi.org/10.1176/appi.ajp.2016.15070854>.
- ⁸² Caine, E.D. (2013). Forging an agenda for suicide prevention in the United States. *Am J Public Health*, 103, 822–829. doi:10.2105/AJPH.2012.301078.
- ⁸³ Bernert, R.A. (2018). Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved on December 14, 2018 from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ⁸⁴ Ibid.
- ⁸⁵ Henderson, C., Evans-Lacko, S., & Thornicroft, G. (2013). Mental illness stigma, help seeking, and public health programs. *American journal of public health*, 103(5), 777–780.
- ⁸⁶ Caine, E.D. (2013). Forging an agenda for suicide prevention in the United States. *Am J Public Health*, 103, 822–829. doi:10.2105/AJPH.2012.301078.
- ⁸⁷ Bernert, R.A. (2018). Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved on December 14, 2018 from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ⁸⁸ Ibid.
- ⁸⁹ Ibid.
- ⁹⁰ Ibid.
- ⁹¹ Ibid.
- ⁹² The Joint Commission. (2016). Detecting and treating suicide ideation in all settings. *Sentinel Event Alert*, 56, 1-7. Retrieved December 19, 2018 from https://www.jointcommission.org/assets/1/18/SEA_56_Suicide.pdf.
- ⁹³ Visit <https://zerosuicide.sprc.org/> for more information on the Zero Suicide Initiative.
- ⁹⁴ Ahmedani, B.K., Simon, G.E., Stewart, C., Beck, A., Waitzfelder, B.E., Rossom, R., et al. (2014). Health care contacts in the year before suicide death. *J Gen Intern Med*, 29(6), 870 – 877.

-
- ⁹⁵ Labouliere, C. D., Vasan, P., Kramer, A., Brown, G., Green, K., Rahman, M., ... Stanley, B. (2018). "Zero Suicide" - A model for reducing suicide in United States behavioral healthcare. *Suicidologi*, 23(1), 22–30.
- ⁹⁶ Coffey, C.E. (2007). Building a system of perfect depression care in behavioral health. *Joint Commission Journal on Quality and Patient Safety*, 33, 193–199.
- ⁹⁷ Bernert, R. A. (2018). Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved December 14, 2018 from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ⁹⁸ James, G., Witten, D., Hastie, T., Tibshivani, R. (Eds). (2013). *An Introduction to Statistical Learning: with Applications in R*. New York: Springer Publishing.
- ⁹⁹ Barak-Corren, Y., Castro, V. M., Javitt, S., Hoffnagle, A. G., Dai, Y., Perlis, R.H., Nack, M. K., Smoller, J. W., Reis, B. Y. (2016). Predicting suicidal behavior from longitudinal electronic health records. *American Journal of Psychiatry*, 174(2), 154–162.
- ¹⁰⁰ Walsh, C.G., Ribeiro, J.D., & Franklin, J.C. (2017). Predicting risk of suicide attempts over time through machine learning. *Clinical Psychological Science*, 5(3), 457–469. <https://doi.org/10.1177/2167702617691560>.
- ¹⁰¹ Ibid.
- ¹⁰² McCarthy, J. F., Bossarte, R. M., Katz, I. R., Thompson, C., Kemp, J., Hannemann, C. M., Nielson, C., & Schoenbaum, M. (2015). Predictive modeling and concentration of the risk of suicide: Implications for preventive interventions in the US Department of Veterans Affairs. *American Journal of Public Health*, 105(9), 1935-1942.
- ¹⁰³ Bernert, R.A. (2018). *Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California*. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ¹⁰⁴ Ibid.
- ¹⁰⁵ United States Food and Drug Administration, United States Department of Health and Human Services. (2012). *Guidance for Industry: Suicidality: Prospective Assessment of Occurrence in Clinical Trials, Draft Guidance*. Retrieved on March 15, 2019 from <http://www.fda.gov/downloads/Drugs/Guidances/UCM225130.pdf>.
- ¹⁰⁶ Bernert, R. A., Hom, M. A., & Roberts, L. W. (2014). A review of multidisciplinary clinical practice guidelines in suicide prevention: Toward an emerging standard in suicide risk assessment and management, training and practice. *Academic Psychiatry*, 38(5), 585-592. <https://doi:10.1007/s40596-014-0180-1>.
- ¹⁰⁷ Caine, E.D. (2013). Forging an agenda for suicide prevention in the United States. *Am J Public Health*, 103, 822–829. doi:10.2105/AJPH.2012.301078.
- ¹⁰⁸ Crosby, A. E., Ortega, L., & Melanson, C. (2011). *Self-directed Violence Surveillance: Uniform Definitions and Recommended Data Elements*, Version 1.0. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control.

-
- ¹⁰⁹ Yip, P. S., Caine, E., Yousuf, S., Chang, S. S., Wu, K. C., & Chen, Y. Y. (2012). Means restriction for suicide prevention. *Lancet (London, England)*, 379(9834), 2393-2399.
- ¹¹⁰ Caine, E. D. (2013). Forging an agenda for suicide prevention in the United States. *The American Journal of Public Health*, 103, 822-829. <https://doi:10.2105/AJPH.2012.301078>.
- ¹¹¹ Bernert, R. A. (2018). Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved December 14, 2018 from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.
- ¹¹² Ibid.
- ¹¹³ Meyer, R.E., Salzman, C., Youngstrom, E.A., Clayton, P.J., Goodwin, F.K., Mann, J.J., Alphas, L.D., Broich, K., Goodman, W.K., Greden, J.F., Meltzer, H.Y., Normand, S.L., Posner, K., Shaffer, D., Oquendo, M.A., Stanley, B., Trivedi, M.H., Turecki, G., Beasley, C.M., Beautrais, A.L., Bridge, J.A., Brown, G.K., Revicki, D.A., Ryan, N.D., & Sheehan, D.V. (2010). Suicidality and risk of suicide: definition, drug safety concerns, and a necessary target for drug development: a consensus statement. *J Clin Psychiatry*, 71(8), e1-e21.
- ¹¹⁴ Miller, M., Azrael, D., & Barber, C. (2012). Suicide mortality in the United States: The importance of attending to method in understanding population-level disparities in the burden of suicide. *Annual Review of Public Health*, 33(1), 393-408.
- ¹¹⁵ California Department of Public Health (2019). *Injury Data Brief: Suicides among Veterans in California, 2017*. Sacramento, CA: California Department of Public Health. Retrieved on March 26, 2019 from <https://www.cdph.ca.gov/Programs/CCDCPHP/DCDIC/SACB/CDPH%20Document%20Library/Violence%20Prevention%20Initiative/CA%20Veteran%20Suicides%202017%20FINALa%203%2011%2019.pdf>.
- ¹¹⁶ Centers for Disease Control, National Center for Health Statistics. *Suicide mortality by state, 2017*. Retrieved March 15, 2019 from <https://www.cdc.gov/nchs/pressroom/sosmap/suicide-mortality/suicide.htm>.
- ¹¹⁷ Anestis, M. D., & Anestis, J. C. (2015). Suicide Rates and State Laws Regulating Access and Exposure to Handguns. *American Journal of Public Health*, 105(10), 2049-2058. doi:10.2105/AJPH.2015.302753.
- ¹¹⁸ Ibid.
- ¹¹⁹ California Department of Public Health (2019). *Preventing Violence in California: Data Brief 1: Overview of Homicide and Suicide Deaths in California*. Sacramento, CA: California Department of Public Health.
- ¹²⁰ Ibid.
- ¹²¹ Miller, M., Barber, C., White, R. A., & Azrael, D. (2013). Firearms and suicide in the United States: is risk independent of underlying suicidal behavior? *Am J Epidemiol.*, 178(6), 946-955.
- ¹²² California Department of Public Health (CDPH) Vital Statistics Death File (2017).
- ¹²³ Ibid.
- ¹²⁴ California Department of Public Health (2019). *Preventing Violence in California: Data Brief 1: Overview of Homicide and Suicide Deaths in California*. Sacramento, CA: California Department of Public Health.

-
- ¹²⁵ Ibid.
- ¹²⁶ Ibid.
- ¹²⁷ Sullivan, E.M., Annest, J.L., Simon, T.R., Luo, F., & Dahlberg, L.L. (2015). Suicide trends among persons aged 10-24 years – United States, 1994-2012. *MMWR Morbidity and Mortality Weekly Report*, 64(8), 201-205.
- ¹²⁸ California Department of Public Health (2019). *Preventing Violence in California: Data Brief 1: Overview of Homicide and Suicide Deaths in California*. Sacramento, CA: California Department of Public Health.
- ¹²⁹ Ibid.
- ¹³⁰ Centers for Disease Control and Prevention. (2017). *WISQARS website and Fatal injury reports*. Retrieved November 28, 2018 from <https://www.cdc.gov/injury/wisqars/fatal.html>.
- ¹³¹ Elnour, A. A., & Harrison, J. (2008) *Lethality of suicide methods*. *Injury Prevention*, 14, 39–45.
- ¹³² Canetto, S.S., & Sakinofsky, I. (1998). The gender paradox in suicide. *Suicide Life Threat Behav*, 28, 1–23.
- ¹³³ Zhang, J., Jiang, C., Jia, S., & Wieczorek, W. F. (2002). An overview of suicide research in China. *Archives of Suicide Research: Official Journal of the International Academy for Suicide Research*, 6(2), 167–184. <https://doi:10.1080/13811110208951174>.
- ¹³⁴ California Department of Public Health (2019). *Preventing Violence in California: Data Brief 1: Overview of Homicide and Suicide Deaths in California*. Sacramento, CA: Author.
- ¹³⁵ Ibid.
- ¹³⁶ Conwell, Y., Duberstein, P. R., Cox, C., et al. (1998). Age differences in behaviors leading to completed suicide. *Am J Geriatr Psychiatry*, 6(2), 122–126.
- ¹³⁷ California Department of Public Health (2019). *Preventing Violence in California: Data Brief 1: Overview of Homicide and Suicide Deaths in California*. Sacramento, CA: Author.
- ¹³⁸ Ibid.
- ¹³⁹ Ibid.
- ¹⁴⁰ Centers for Disease Control and Prevention. (2017). *WISQARS website and fatal injury reports*. Retrieved November 28, 2018 from <https://www.cdc.gov/injury/wisqars/fatal.html>.
- ¹⁴¹ California Department of Public Health (2019). *Injury Data Brief: Suicides among Veterans in California, 2017*. Sacramento, CA: Author. Retrieved March 26, 2019 from <https://www.cdph.ca.gov/Programs/CCDPHP/DCDIC/SACB/CDPH%20Document%20Library/Violence%20Prevention%20Initiative/CA%20Veteran%20Suicides%202017%20FINALa%203%2011%2019.pdf>.
- ¹⁴² Ibid.
- ¹⁴³ Ibid.
- ¹⁴⁴ Ibid.
- ¹⁴⁵ Barber, C. W., & Miller, M. J. (2014). Reducing a suicidal person’s access to lethal means of suicide: A research agenda. *American Journal of Preventative Medicine*, 47(3), S264-S272.
- ¹⁴⁶ California Department of Justice. *Death in Custody and Arrest-Related Deaths*. Database available by visiting <https://openjustice.doj.ca.gov/data>. Data mandated per Government Code Section 12525.
- ¹⁴⁷ California Department of Justice. *Death in Custody Context*. Retrieved December 13, 2018 from <https://openjustice.doj.ca.gov/data>.
- ¹⁴⁸ Ibid.

¹⁴⁹ Ibid.

¹⁵⁰ Office of Statewide Health Planning and Development (OSHPD). *Hospital Emergency Department - Characteristics by Facility, 2017*. Retrieved September 28, 2018 from <https://data.chhs.ca.gov/dataset/hospital-emergency-department-characteristics-by-facility-pivot-profile>.

¹⁵¹ Substance Abuse and Mental Health Services Administration. *Behavioral Health Barometer: California, Volume 4: Indicators as measured through the 2015 National Survey on Drug Use and Health, the National Survey of Substance Abuse Treatment Services, and the Uniform Reporting System*. HHS Publication No. SMA-17-Baro-16-States-CA. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2017.

¹⁵² Data Source: WestEd, California Healthy Kids Survey. California Department of Education (Jul. 2017). As cited on kidsdata.org, a program of the Lucile Packard Foundation for Children's Health.

¹⁵³ Stone, D. M., Holland, K. M., Bartholow, B., E Logan, J., LiKamWa McIntosh, W., Trudeau, A., & Rockett, I. (2017). Deciphering Suicide and Other Manners of Death Associated with Drug Intoxication: A Centers for Disease Control and Prevention Consultation Meeting Summary. *American journal of public health, 107*(8), 1233–1239. doi:10.2105/AJPH.2017.303863.

¹⁵⁴ California Government Code Section 27491.

¹⁵⁵ U.S. Department of Justice, Office of Justice Programs, National Institute of Justice. (2011). *Death investigations: A guide for the scene investigator – technical update*. Retrieved on December 11, 2018 from <https://www.ncjrs.gov/pdffiles1/nij/234457.pdf>.

¹⁵⁶ Stone, D. M., Holland, K. M., Bartholow, B., E Logan, J., LiKamWa McIntosh, W., Trudeau, A., & Rockett, I. (2017). Deciphering Suicide and Other Manners of Death Associated with Drug Intoxication: A Centers for Disease Control and Prevention Consultation Meeting Summary. *American journal of public health, 107*(8), 1233–1239. doi:10.2105/AJPH.2017.303863.

¹⁵⁷ Mohler, B., & Earls, F. (2001). Trends in adolescent suicide: misclassification bias? *American Journal of Public Health, 91*(1), 150–153.

¹⁵⁸ Bernert, R. A. (2018). *Emerging best practices and innovations in suicide prevention: Toward an updated strategic plan for California*. Sacramento, CA: Mental Health Services Oversight and Accountability Commission. Retrieved from http://mhsoac.ca.gov/sites/default/files/documents/2018-11/Policy%20Brief_Emerging%20best%20practices%20in%20suicide%20prevention_10.17.2018.pdf.

¹⁵⁹ Ibid.

¹⁶⁰ Ibid.

¹⁶¹ Ibid.

¹⁶² Miller, M., Azrael, D., & Barber, C. (2012). Suicide mortality in the United States: The importance to attending to method in understanding population-level disparities in the burden of suicide. *Annual Review of Public Health, 33*, 393–408.

¹⁶³ Gould, M. S. (1990). *Suicide clusters and media exposure*. In: Blumenthal S, Kupfer D, editors. *Suicide over the life cycle: Risk factors, assessment, and treatment of suicidal patients*. Washington, DC: American Psychiatric Association; pp. 517–532.

¹⁶⁴ Suicide Prevention Resource Center, & Rodgers, P. (2011). *Understanding risk and protective*

factors for suicide: A primer for preventing suicide. Newton, MA: Education Development Center, Inc.

¹⁶⁵ World Health Organization. (2012). *Public health action for the prevention of suicide: A framework*. Geneva, Switzerland: Author.

¹⁶⁶ Suicide Prevention Resource Center, & Rodgers, P. (2011). *Understanding risk and protective factors for suicide: A primer for preventing suicide*. Newton, MA: Education Development Center, Inc.

¹⁶⁷ Ibid.

¹⁶⁸ Colucci, E. & Martin, G. (2008). Religion and spirituality along the suicidal path. *Suicide & Life-Threatening Behavior*, 38(2), 229–244.

¹⁶⁹ Lun, V. M. C., & Bond, M. H. (2013). Examining the relation of religion and spirituality to subjective well-being across national cultures. *Psychology of Religion and Spirituality*, 5, 304–315.

¹⁷⁰ Goldston, D. B., Molock, S. D., Whitbeck, L. B., Murakami, J. L., Zayas, L. H., & Hall, G. C. (2008). Cultural considerations in adolescent suicide prevention and psychosocial treatment. *The American Psychologist*, 63, 14–31.

¹⁷¹ Wachholtz, A., & Sambamoorthi, U. (2011). National trends in prayer use as a coping mechanism for health concerns: Changes from 2002 to 2007. *Psychology of Religion and Spirituality*, 3, 67–77.

¹⁷² Lytle, M.C., Blossnich, J.R., De Luca, S.M., & Brownson, C. (2018). Association of religiosity with sexual minority suicide ideation and attempt. *American Journal of Preventative Medicine*, 54(5), 644–651. DOI: <https://doi.org/10.1016/j.amepre.2018.01.019>.

¹⁷³ Suicide Prevention Resource Center (n.d.). Risk and Protective Factors retrieved from <https://www.sprc.org/about-suicide/risk-protective-factors> and Warning Signs retrieved from <https://www.sprc.org/about-suicide/warning-signs>.

¹⁷⁴ Ibid.

¹⁷⁵ Ibid.

¹⁷⁶ Bertolote, J. M., & Fleischmann, A. (2002). Suicide and psychiatric diagnosis: A worldwide perspective. *World Psychiatry*, 1, 181–185.

¹⁷⁷ Centers for Disease Control and Prevention (2013). Suicide among adults aged 35–64 years: United States, 1999–2010. *MMWR Morb Mortal Wkly Rep*, 62, 321–325.

¹⁷⁸ Ibid.

¹⁷⁹ Drapeau, C. W., & McIntosh, J. L. (for the American Association of Suicidology). (2018). *U.S.A. suicide 2017: Official final data*. Washington, DC: American Association of Suicidology, dated December 10, 2018, downloaded from <http://www.suicidology.org>.

¹⁸⁰ Ibid.

¹⁸¹ Ibid.

¹⁸² Conwell, Y. (2014). Suicide later in life: Challenges and priorities for prevention. *American Journal of Preventive Medicine* 47(3 Suppl. 2), S244–S250.

¹⁸³ Cooper, J., Kapur, N., Webb, R., Lawlor, M., Guthrie, E., Mackway-Jones, K., & Appleby, L. (2005). Suicide after deliberate self-harm: A 4-year cohort study. *The American Journal of Psychiatry*, 162, 297–303.

¹⁸⁴ Qin, P., & Nordentoft, M. (2005). Suicide risk in relation to psychiatric hospitalization: Evidence based on longitudinal registers. *Archives of General Psychiatry*, 62(4), 427–432.

-
- ¹⁸⁵ Valenstein, M., Kim, H.M., & Ganoczy, D., et al. (2009). Higher-risk periods for suicide among VA patients receiving depression treatment: prioritizing suicide prevention efforts. *Journal of Affective Disorders*, 112(1-3), 50–58.
- ¹⁸⁶ Department of Veterans Affairs, Veterans Health Administration, Office of Mental Health and Suicide Prevention. (2018). *Veteran Suicide Data Report, 2005–2016*. Retrieved from https://www.mentalhealth.va.gov/docs/data-sheets/OMHSP_National_Suicide_Data_Report_2005-2016_508-compliant.pdf.
- ¹⁸⁷ Ibid.
- ¹⁸⁸ Ibid.
- ¹⁸⁹ Barber, C. W., & Miller, M. J. (2014). Reducing a suicidal person's access to lethal means of suicide: A research agenda. *American Journal of Preventative Medicine*, 47(3), S264-S272.
- ¹⁹⁰ U.S. Department of Veterans Affairs. (2016). *National Strategy for Preventing Veteran Suicide (2018-2028)*. Retrieved May 31, 2019 from https://www.mentalhealth.va.gov/suicide_prevention/docs/Office-of-Mental-Health-and-Suicide-Prevention-National-Strategy-for-Preventing-Veterans-Suicide.pdf.
- ¹⁹¹ Ibid.
- ¹⁹² Ibid.
- ¹⁹³ Demers, A. (2011). When veterans return: The role of community in reintegration. *Journal of Loss and Trauma*, 16, 160–179.
- ¹⁹⁴ World Health Organization (2016). FAQ on health and sexual diversity - An introduction to key concepts. Geneva: Author; Licence: CC BY-NC-SA 3.0 IGO. Retrieved May 2, 2019 from <https://www.who.int/gender-equity-rights/news/20170329-health-and-sexual-diversity-faq.pdf?ua=1>.
- ¹⁹⁵ Zhao, Y., Montoro, R., Igartua, K., & Thombs, B. D. (2010). Suicidal ideation and attempt among adolescents reporting “unsure” sexual identity or heterosexual identity plus same-sex attraction or behavior: Forgotten groups? *Journal of the American Academy of Child and Adolescent Psychiatry*, 49(2), 104–113.
- ¹⁹⁶ Hottes, T. S., Bogaert, L., Rhodes, A. E., Brennan, D. J., & Gesink, D. (2016). Lifetime prevalence of suicide attempts among sexual minority adults by study sample strategies: A systematic review and meta-analysis. *American Journal of Public Health*, 106(5), e1-12. <https://doi:10.2105/AJPH.2016.303088>.
- ¹⁹⁷ Salway, T., Ross, L. E., Fehr, C. P., Burley, J., Asadi, S., Hawkins, B., Tarasoff, L. A. (2018). A systematic review and meta-analysis of disparities in the prevalence of suicide ideation and attempt among bisexual populations. *Archives of Sexual Behavior*. <https://doi.org/10.1007/s10508-018-1150-6>.
- ¹⁹⁸ Shearer, A., Herres, J., Kodish, T., Squitieri, H., James, K., Russon, J., Atte, T., & Diamond, G. S. (2016). Differences in mental health symptoms across lesbian, gay, bisexual, and questioning youth in primary care settings. *Journal of Adolescent Health*, 59(1), 38 – 43.
- ¹⁹⁹ Ibid.
- ²⁰⁰ Haas, A. P., Eliason, M., Mays, V. M., Mathy, R. M., Cochran, S. D., D'Augelli, A. R., ... Clayton, P. J. (2011). Suicide and suicide risk in lesbian, gay, bisexual, and transgender populations: review and recommendations. *Journal of homosexuality*, 58(1), 10–51.

-
- ²⁰¹ James, S. E., Herman, J. L., Rankin, S., Keisling, M., Mottet, L., & Anafi, M. (2016). *The Report of the 2015 U.S. Transgender Survey*. Washington, DC: National Center for Transgender Equality.
- ²⁰² Haas, A., Rodgers, P., & Herman, J.L. (2014). *Suicide attempts among transgender and gender non-conforming adults: Findings of the National Transgender Discrimination Survey*. Retrieved January 28, 2019 from <https://williamsinstitute.law.ucla.edu/wp-content/uploads/AFSP-Williams-Suicide-Report-Final.pdf>.
- ²⁰³ Ryan, C., Huebner, D., Diaz, R., & Sanchez, J. (2009). Family rejection as a predictor of negative health outcomes in White and Latino LGB young adults. *Pediatrics.*, 123, 346–352.
- ²⁰⁴ U.S. Department of Health and Human Services, Office of Minority Health. (n.d.) *Suicide and Suicide Prevention 101*. Retrieved from <http://minorityhealth.hhs.gov/templates/browse.aspx?lvl=3&lvlid=136>.
- ²⁰⁵ National Center for Injury Prevention and Control, CDC. (2016). NCHS Vital Statistics System for numbers of deaths. WISQARS: Web-based Injury Statistics Query and Reporting System. Retrieved September 12, 2018 from <https://webappa.cdc.gov>.
- ²⁰⁶ Bridge, J. A., Horowitz, L. M., Fontanella, C. A., Sheftall, A., Greenhouse, J., Kelleher, K. J., & Campo, J. V. (2018). Age-Related racial disparity in suicide rates among US youths from 2001 through 2015. *JAMA Pediatrics*, 172(7), 697–699.
- ²⁰⁷ Crosby, A. E., Han, B., Ortega, L. A. G., Parks, S. E., & Gfoerer, J. (2011). Suicidal thoughts and behaviors among adults aged ≥ 18 years - United States, 2008-2009. *MMWR Surveillance Summaries*, 60 (SS-13), 1-22. Retrieved from www.cdc.gov/mmwr/preview/mmwrhtml/ss6013a1.htm?s_cid=ss6013a1_e.
- ²⁰⁸ Chu, J. P., Goldblum, P., Floyd, R., & Bongar, B. (2010). The cultural theory and model of suicide. *Applied and Preventative Psychology*, 14(1-4), 25-40.
- ²⁰⁹ U.S. Centers for Disease Control and Prevention. (2008). Youth Risk Behavior Surveillance—Selected Steps Communities, 2007. *Morbidity and Mortality Weekly Reports*, 57(SS–12), 1–27.
- ²¹⁰ Zayas, L. H. (2011). *Latinas attempting suicide: When cultures, families, and daughters collide*. New York, NY, US: Oxford University Press.
- ²¹¹ National Advisory Committee on Rural Health and Human Services (2017). *Understanding the impact of suicide in rural America: Policy brief and recommendations*. Retrieved February 12, 2019 <https://www.hrsa.gov/sites/default/files/hrsa/advisory-committees/rural/publications/2017-impact-of-suicide.pdf>.
- ²¹² Ibid.
- ²¹³ Ibid.
- ²¹⁴ Ibid.
- ²¹⁵ Peterson, C., Stone, D. M., & Marsh, S. M., et al. (2018). Suicide Rates by Major Occupational Group — 17 States, 2012 and 2015. *MMWR Morb Mortal Wkly Rep*, 67, 1253–1260. <http://dx.doi.org/10.15585/mmwr.mm6745a1>.
- ²¹⁶ McIntosh, W. L., Spies, E., Stone, D. M., Lokey, C. N., Trudeau, A. T., Bartholow, B. (2016). Suicide Rates by Occupational Group — 17 States, 2012. *MMWR Morb Mortal Wkly Rep*, 65, 641–645.
- ²¹⁷ Ibid.

-
- ²¹⁸ Stanley, I. H., Hom, M. A., & Joiner, T. E. (2016). A systematic review of suicidal thoughts and behaviors among police officers, firefighters, EMTs, and paramedics. *Clinical Psychology Review*, 44, 25–44. <https://doi.org/10.1016/j.cpr.2015.12.002>.
- ²¹⁹ Hayes, L. (2006). *Suicide prevention in correctional facilities: An overview*. In M. Puisis (Ed.), *Clinical Practice in Correctional Medicine* (pp. 317-340). Philadelphia, PA: Mosby Elsevier.
- ²²⁰ Jenkins, R., Bhugra, D., Meltzer, H., Singleton, N., Bebbington, P., Brugha, T., Coid, J., Farrell, M., Lewis, G., & Paton, J. (2005). Psychiatric and social aspects of suicidal behaviour in prisons. *Psychological Medicine*, 35, 257-269.
- ²²¹ Stack, S. J., & Tsoudis, O. (1997). Suicide Risk among Correctional Officers: A Logistic Regression Analysis. *Archives of Suicide Research*, 3(3), 183-186.
- ²²² Brower, J. (2013). *Review and Input of Correctional Officer Wellness & Safety Literature Review*. OJP Diagnostic Center. Office of Justice Programs.
- ²²³ Mrazek, P. J., & Haggerty, R. J. (1994). *Reducing risks for mental disorders: frontiers for preventive intervention research*. Washington, D.C.: National Academy Press.
- ²²⁴ Ibid.
- ²²⁵ Ibid.
- ²²⁶ Ibid.
- ²²⁷ Cramer, R. J., & Kapusta, N. D. (2017). A Social-Ecological Framework of Theory, assessment, and prevention of suicide. *Frontiers in Psychology*, 8, 1756. <https://doi:10.3389/fpsyg.2017.01756>.
- ²²⁸ Knapp, M., McDaid, D., & Parsonage, M. (2011). *Mental health promotion and mental illness prevention: the economic case*. Department of Health, London, UK. Retrieved January 4, 2019 from [http://eprints.lse.ac.uk/39303/1/Mental_health_promotion_and_mental_illness_prevention\(a_uthor\).pdf](http://eprints.lse.ac.uk/39303/1/Mental_health_promotion_and_mental_illness_prevention(a_uthor).pdf).
- ²²⁹ Lavinghouze, S. R., Snyder, K., & Rieker, P. P. (2014). The component model of infrastructure: a practical approach to understanding public health program infrastructure. *American journal of public health*, 104(8), e14-24.
- ²³⁰ Tobacco Institute. (2009). *Overview of state ASSIST programs*. Bates no. TI25390805. Retrieved January 4, 2019 from <http://legacy.library.ucsf.edu/tid/qlr45b00>.
- ²³¹ Barber, C. W., & Miller, M. J. (2014). Reducing a suicidal person's access to lethal means of suicide: A research agenda. *American Journal of Preventative Medicine*, 47(3), S264-S272.
- ²³² Ibid.
- ²³³ Kreitman, N. (1976). The coal gas story. United Kingdom suicide rates, 1960-71. *British Journal of Preventive & Social Medicine*, 30(2), 86-93.
- ²³⁴ Lubin, G., Werbeloff, N., Halperin, D., Shmushkevitch, M., Weiser, M., & Knobler, H. Y. (2010). Decrease in suicide rates after a change of policy reducing access to firearms in adolescents: a naturalistic epidemiological study. *Suicide and Life-Threatening Behavior*, 40(5), 421–424. <https://doi.org/10.1521/suli.2010.40.5.421>.
- ²³⁵ Knipe, D. W., Chang, S. S., Dawson, A., Eddleston, M., Konradsen, F., Metcalfe, C., & Gunnell, D. (2017). Suicide prevention through means restriction: Impact of the 2008-2011 pesticide restrictions on suicide in Sri Lanka. *PLoS one*, 12(3), e0172893. <https://doi:10.1371/journal.pone.0172893>.

-
- ²³⁶ Shelef, M. (1994). Unanticipated benefits of automotive emission control: Reduction in fatalities by motor vehicle exhaust gas. *Sci. Total Environ.*, 146-147, 93–101. [https://doi:10.1016/0048-9697\(94\)90224-0](https://doi:10.1016/0048-9697(94)90224-0).
- ²³⁷ Mann, J. J., Apter, A., Bertolote, J., Beautrais, A., Currier, D., Haas, A., Hegerl, U., Lonnqvist, J., Malone, K., Marusic, A., Mehlum, L., Patton, G., Phillips, M., Rutz, W., Rhimer, Z., Schmidtke, A., Shaffer, D., Silverman, M., Takahasashi, Y., Varnik, A., Wasserman, D., Wip, P., & Hardin, H. (2005). Suicide prevention strategies: A systematic review. *The Journal of the American Medical Association*, 294, 2064–2074.
- ²³⁸ Beautrais, A. L. (2001). Effectiveness of barriers at suicide jumping sites: A case study. *Australian and New Zealand Journal of Psychiatry*, 35, 557–562.
- ²³⁹ Beautrais, A. L., Gibb, S. J., Fergusson, D. M., Horwood, L. J., & Larkin, G. L. (2009). Removing bridge barriers stimulates suicides: an unfortunate natural experiment. *Australian and New Zealand Journal of Psychiatry*, 43(6), 495–497.
- ²⁴⁰ Law, C. K., Sveticic, J., & De Leo, D. (2014). Restricting access to a suicide hotspot does not shift the problem to another location. An experiment of two river bridges in Brisbane, Australia. *Australian and New Zealand Journal of Public Health*, 38(2), 134–138.
- ²⁴¹ Surface Transportation Block Grant Program (STBG). Federal Highway Administration (FHWA). (2016). *CA GOVERNMENT CODE, TITLE 2, CHAPTER 2, SEC 14527.1 and 23 USC 133(b)(7). The STBG promotes flexibility in State and local transportation funding decisions to best address State and local transportation needs. FAST Act § 1109(a)*.
- ²⁴² Sarchiapone, M., Mandelli, L., Iosue, M., Andrisano, C., & Roy, A. (2011). Controlling access to suicide means. *International Journal of Environmental Research and Public Health*, 8(12), 4550–62.
- ²⁴³ Draper, J. (2008). Suicide prevention on bridges: The National Suicide Prevention Lifeline position. Retrieved on February 12, 2019 from https://suicidepreventionlifeline.org/wp-content/uploads/2016/09/SUICIDE_BRIDGES_Lifeline_Position_Paper_Final_6-16-08.pdf.
- ²⁴⁴ Anglemeyer, A., Horvath, T., & Rutherford, G. (2014). The accessibility of firearms and risk for suicide and homicide victimization among household Members: A systematic review and meta-analysis. *Annals of Internal Medicine*, 160(2), 101–110. <https://doi:10.7326/M13-1301>.
- ²⁴⁵ Kellermann, A. L., Rivara, F. P., Somes, G., Reay, D. T., Francisco, J., Banton, J. G., Prodzinski, J., Flighter, C., & Hackman, B. B. (1992). Suicide in the home in relation to gun ownership. *New England Journal of Medicine*, 327, 467–472.
- ²⁴⁶ Siegel, M., Pahn, M., Xuan, Z., Fleegler, E., & Hemenway, D. (2019). The impact of state firearm laws on homicide and suicide deaths in the USA, 1991–2016: a panel study. *Journal of General Internal Medicine*. <https://doi:10.1007/s11606-019-04922-x>.
- ²⁴⁷ Washington State Legislature. (2014). Washington State *RCW 9.41.113: Firearm sales or transfers—Background checks—Requirements—Exceptions*. Washington: Washington State Legislature.
- ²⁴⁸ Kivisto, A. J., & Phalen, P. L. (2018). Effects of risk-based firearm seizure laws in Connecticut and Indiana on suicide rates, 1981–2015. *Psychiatric Services*, 69(8), 855–862. <https://doi.org/10.1176/appi.ps.201700250>.
- ²⁴⁹ American Psychiatric Association, Ad Hoc Workgroup of the Council on Psychiatry and Law. (2018). *Resource document on risk-based gun removal laws*. Retrieved November 8, 2018 from

<https://www.psychiatry.org/psychiatrists/search-directories-databases/library-and-archive/resource-documents>.

²⁵⁰ California Penal Code Sections 18100 to 18205.

²⁵¹ Superior California Suicide Prevention Network. (2014). *Toolkit for Communities Collaborating in Suicide Prevention and Firearms Safety*. Retrieved December 28, 2018 from <https://www.co.shasta.ca.us/docs/libraries/hhsa-docs/mental-wellness/FirearmToolkit.pdf>.

²⁵² Information retrieved on March 15, 2019 from <https://afsp.org/american-foundation-suicide-prevention-national-shooting-sports-foundation-partner-help-prevent-suicide/>.

²⁵³ Spicer, R.S., & Miller, T.R. (2000). Suicide acts in 8 states: Incidence and case fatality rates by demographics and method. *American Journal of Public Health*, 90(12), 1885-1891.

²⁵⁴ California Civil Code Section 1714.22

²⁵⁵ Bryan, C. J., Stone, S. L., & Rudd, M. D. (2011). A practical, evidence-based approach for means-restriction counseling with suicidal patients. *Professional Psychology: Research and Practice*, 42(5), 339-346.

²⁵⁶ Kruesi, M. J., Grossman, J., Pennington, J. M., Woodward, P. J., Duda, D., & Hirsch, J. G. (1999). Suicide and violence prevention: parent education in the emergency department. *Journal of the American Academy of Child & Adolescent Psychiatry*, 38(3), 250-255.

²⁵⁷ Betz, M. E., Miller, M., Barber, C., Beaty, B., Miller, I., Camargo, C. A., & Boudreaux, E. (2016). Lethal means access and assessment among suicidal emergency department patients. *Depression and Anxiety*, 33(6), 502-511.

²⁵⁸ Visit the Suicide Prevention Resource Center at <https://training.sprc.org/>.

²⁵⁹ Johnson, R. M., Frank, E. M., Ciocca, M., & Barber, C. W. (2011) Training mental healthcare providers to reduce at-risk patients' access to lethal means of suicide: evaluation of the CALM Project. *Archives of Suicide Research*, 15(3), 259-264.

²⁶⁰ Centers for Disease Control and Prevention (2009). *Strategic direction for the prevention of suicidal behavior: promoting individual, family, and community connectedness to prevent suicidal behavior*. Retrieved from http://www.cdc.gov/ViolencePrevention/pdf/Suicide_Strategic_Direction_Full_version-a.pdf.

²⁶¹ Greden, J. F., Valenstein, M., Spinner, J., Blow, A., Gorman, L. A., Dalack, G. W., Marcus, S., & Kees, M. (2010). Buddy-to-Buddy, a citizen soldier peer support program to counteract stigma, PTSD, depression, and suicide. *Annals of the New York Academy of Sciences*, 1208(1), 90-97. <https://doi.org/10.1111/j.1749-6632.2010.05719.x>.

²⁶² Whitlock, J., Wyman, P. A., & Moore, S. R. (2014). Connectedness and suicide prevention in adolescents: Pathways and implications. *Suicide and Life-Threatening Behavior*, 44, 246-272.

²⁶³ Saewyc, E., Konishi, C., Rose, H., & Homma, Y. (2014). School-based strategies to reduce suicidal ideation, suicide attempts, and discrimination among sexual minority and heterosexual adolescents in Western Canada. *International Journal of Child Youth and Family Studies*, 5(1), 89-112. <https://doi.org/10.18357/ijcyfs.saewyce.512014>.

²⁶⁴ Ryan, C., Huebner, D., Diaz, R., & Sanchez, J. (2009). Family rejection as a predictor of negative health outcomes in White and Latino LGB young adults. *Pediatrics*, 123, 346-352.

²⁶⁵ Centers for Disease Control and Prevention (2009). *Strategic direction for the prevention of suicidal behavior: promoting individual, family, and community connectedness to prevent suicidal behavior*. Retrieved from http://www.cdc.gov/ViolencePrevention/pdf/Suicide_Strategic_Direction_Full_version-a.pdf.

-
- ²⁶⁶ Lester, L., & Cross, D. (2015). The relationship between school climate and mental and emotional wellbeing over the transition from primary to secondary school. *Psychology of Well-Being*, 5(1), 9.
- ²⁶⁷ Wilkins, N., Tsao, B., Hertz, M., Davis, R., & Klevens, J. (2014). *Connecting the Dots: An Overview of the Links Among Multiple Forms of Violence*. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention and Oakland, CA: Prevention Institute.
- ²⁶⁸ Ibid.
- ²⁶⁹ Ibid.
- ²⁷⁰ Wilkins, N., Myers, L., Kuehl, T., Bauman, A., & Hertz, M. (2018). Connecting the dots: State health department approaches to addressing shared risk and protective factors across multiple forms of violence. *Journal of Public Health Management and Practice: JPHMP*, 24(Suppl 1 INJURY AND VIOLENCE PREVENTION), S32–S41.
<http://doi.org/10.1097/PHH.0000000000000669>.
- ²⁷¹ Ibid.
- ²⁷² Joe, S., Canetto, S. S., & Romer, D. (2008). Advancing prevention research on the role of culture in suicide prevention. *Suicide & Life-threatening Behavior*, 38(3), 354–362.
- ²⁷³ Kellam, S. G., Mackenzie, A. C. L., Brown, C. H., Poduska, J. M., Wang, W., Petras, H., & Wilcox, H. C. (2011). The Good Behavior Game and the future of prevention and treatment. *Addiction Science and Clinical Practice*, 6, 73–84.
- ²⁷⁴ LaFromboise, T., & Howard-Pitney, B. (1995). The Zuni life skills development curriculum: Description and evaluation of a suicide prevention program. *Journal of Counseling Psychology*, 42(4), 479.
- ²⁷⁵ De Leo, D., & Heller, T. (2008). Social modeling in the transmission of suicidality. *Crisis: The Journal of Crisis Intervention and Suicide Prevention*, 29, 11–19.
- ²⁷⁶ Gould, M. S. (1990). *Suicide clusters and media exposure*. In: In: Blumenthal S, Kupfer D, editors. *Suicide over the life cycle: Risk factors, assessment, and treatment of suicidal patients*. Washington, DC: American Psychiatric Association; pp. 517–532.
- ²⁷⁷ Hazell, P. (1993). Adolescent suicide clusters—evidence, mechanisms and prevention. *Australian and New Zealand Journal of Psychiatry*, 4(27), 653–665.
- ²⁷⁸ Gould, M. S., Kleinman, M., Lake, A., Forman, J., & Midle, J. (2014). Newspaper coverage of suicide and initiation of suicide clusters in teenagers in the USA, 1988–96: A retrospective, population-based, case control study. *The Lancet Psychiatry*, 1(1), 34–43.
- ²⁷⁹ Gould, M. S. (2001). Suicide and the media. *Annals of the New York Academy of Sciences*, 932, 200–221.
- ²⁸⁰ Williams, J. (2011). The effect on young people of suicide reports in the media. *Mental Health Practice*, 8(14), 34–36.
- ²⁸¹ Thomas, K., Chang, S. S., & Gunnell, D. (2011). Suicide epidemics: the impact of newly emerging methods on overall suicide rates—a time trends study. *BMC Public Health*, 11, 314.
- ²⁸² SAMHSA, U.S. Department of Health and Human Services, Entertainment Industries Council, Inc., ENCORE Management Corporation. (n.d.). *Picture This: Depression and Suicide Prevention* (contract number 280-02-07010). Entertainment Industries Council, Inc.
- ²⁸³ Ibid.

²⁸⁴ Ibid.

²⁸⁵ While, D., Bickley, H., Roscoe, A., et al. (2012). Implementation of mental health service recommendations in England and Wales and suicide rates, 1997–2006: a cross-sectional and before-and-after observational study. *Lancet.*, 379(9820), 1005–1012.

²⁸⁶ Bashshur, R. L., Shannon, G. W., Bashshur, N., & Yellowlees, P. M. (2016). The Empirical evidence for telemedicine interventions in mental disorders. *Telemedicine Journal and E-Health*, 22(2), 87–113.

²⁸⁷ Mohr, D. C., Ho, J., Duffecy, J., et al. (2012). Effect of telephone-administered vs face-to-face cognitive behavioral therapy on adherence to therapy and depression outcomes among primary care patients: a randomized trial. *Journal of the American Medicine Association*, 307(21), 2278–2285.

²⁸⁸ Gilmore, A. K., & Ward-Ciesielski, E. F. (2017). Perceived risks and use of psychotherapy via telemedicine for patients at risk for suicide. *Journal of Telemedicine and Telecare*, 1357633X17735559. Advance online publication.doi:10.1177/1357633X17735559.

²⁸⁹ Pickering, T. A., Wyman, P. A., Schmeelk-Cone, K., Hartley, C., Valente, T. W., Pisani, A. R., Rullison, K., Brown, C., & LoMurray, M. (2018). Diffusion of a peer-led suicide preventive intervention through school-based student peer and adult networks. *Frontiers in Psychiatry*, 9, 598. <https://doi:10.3389/fpsy.2018.00598>.

²⁹⁰ Ibid.

²⁹¹ Hanisch, S. E., Twomey, C. D., Szeto, A. C., Birner, U. W., Nowak, D., & Sabariego, C. (2016). The effectiveness of interventions targeting the stigma of mental illness at the workplace: a systematic review. *BMC psychiatry*, 16, 1. <https://doi:10.1186/s12888-015-0706-4>.

²⁹² Archer, J., Bower, P., Gilbody, S., Lovell, K., Richards, D., Gask, L., Dickens, C., & Coventry, P. (2012). Collaborative care for depression and anxiety problems. *Cochrane Database of Systematic Reviews* 2012, 10, Art. No.: CD006525. <https://doi:10.1002/14651858.CD006525.pub2>.

²⁹³ American Psychiatric Association and Academy of Psychosomatic Medicine (2016). Dissemination of integrated care within adult primary care settings: The Collaborative Care Model. Retrieved February 12, 2019 from <https://www.psychiatry.org/psychiatrists/practice/professional-interests/integrated-care/learn>.

²⁹⁴ Alexopoulos, G. S., Reynolds, C. F., Bruce, M. L., Katz, I. R., Raue, P. J., Mulsant, B. H., Oslin, D. W., Ten Have, T., PROSPECT Group (2009). Reducing suicidal ideation and depression in older primary care patients: 24-month outcomes of the PROSPECT study. *The American Journal of Psychiatry*, 166(8), 882–890.

²⁹⁵ Unützer, J., Tang, L., Oishi, S., Katon, W., Williams, J., Hunkeler E., & the IMPACT Investigators (2006). Reducing suicidal ideation in depressed older primary care patients. *Journal of the American Geriatrics Society*, 54, 1550–1556.

²⁹⁶ Luoma, J. B., Martin, C. E., & Pearson, J. L. (2002). Contact with mental health and primary care providers before suicide: a review of the evidence. *American Journal of Psychiatry*, 159, 909–916.

²⁹⁷ Lopez, A. D., Mathers, C. D., Ezzati, M., Jamison, D. T., & Murray, C. J. (2006). Global and regional burden of disease and risk factors: Systematic analysis of population health data. *The Lancet*, 367, 1747–1757.

-
- ²⁹⁸ Simon, G. E., Von Korff, M., & Piccinelli, M., et al. (1999). An international study of the relation between somatic symptoms and depression. *The New England Journal of Medicine*, 341, 658–659.
- ²⁹⁹ Mann, J. J., Apter, A., Bertolote, J., Beautrais, A., Currier, D., Haas, A., Hegerl, U., Lonnqvist, J., Malone, K., Marusic, A., Mehlum, L., Patton, G., Phillips, M., Rutz, W., Rhimer, Z., Schmidtke, A., Shaffer, D., Silverman, M., Takahasashi, Y., Varnik, A., Wasserman, D., Wip, P., & Hardin, H. (2005). Suicide prevention strategies: A systematic review. *The Journal of the American Medical Association*, 294, 2064–2074.
- ³⁰⁰ Burnette, C., Ramchand, R., & Ayer, L. (2015). Gatekeeper training for suicide prevention: A theoretical model and review of the empirical literature. *Rand Health Quarterly*, 5(1), 16.
- ³⁰¹ Ibid.
- ³⁰² Ibid.
- ³⁰³ Ibid.
- ³⁰⁴ Ibid.
- ³⁰⁵ Substance Abuse and Mental Health Services Administration. (2014). *Crisis Services: Effectiveness, Cost Effectiveness, and Funding Strategies*. HHS Publication No. (SMA)-14-4848. Rockville, MD: Author.
- ³⁰⁶ National Association of State Mental Health Program Directors (2018). *A comprehensive crisis system: Ending unnecessary emergency room admissions and jail bookings associated with mental illness*. Alexandria, VA: Broadway, E., Covington, D., National Association of State Mental Health Program Directors. Retrieved February 12, 2019 from https://www.nasmhpd.org/sites/default/files/TACPaper5_ComprehensiveCrisisSystem_508C.pdf.
- ³⁰⁷ Substance Abuse and Mental Health Services Administration (2014). *Crisis Services: Effectiveness, Cost Effectiveness, and Funding Strategies*. HHS Publication No. (SMA)-14-4848. Rockville, MD: Author.
- ³⁰⁸ Ibid.
- ³⁰⁹ Acosta, J., Ramchand, R., Jaycox, L.H., Becker, A., & Eberhart, N.K. (2012). Interventions to Prevent Suicide: A Literature Review to Guide Evaluation of California's Mental Health Prevention and Early Intervention Initiative, Santa Monica, Calif.: RAND Corporation, TR-1317-CMHSA, 2012. Retrieved October 18, 2018 from http://www.rand.org/pubs/technical_reports/TR1317.html.
- ³¹⁰ Gould, M. S., Cross, W., Pisani, A. R., Munfakh, J. L., & Kleinman, M. (2013). Impact of Applied Suicide Intervention Skills Training on the National Suicide Prevention Lifeline. *Suicide & Life-threatening Behavior*, 43(6), 676-691.
- ³¹¹ Bernert, R. A., Hom, M. A., & Roberts, L. W. (2014). A review of multidisciplinary clinical practice guidelines in suicide prevention: Toward an emerging standard in suicide risk assessment and management, training and practice. *Academic Psychiatry*, 38(5), 585-592. <https://doi:10.1007/s40596-014-0180-1>.
- ³¹² Jobes, D. A., Gregorian, M. J., & Colborn, V. A. (2018). A stepped care approach to clinical suicide prevention. *Psychological Services*, 15(3), 243-250. <http://dx.doi.org/10.1037/ser0000229>.
- ³¹³ Ibid.
- ³¹⁴ Ibid.

-
- ³¹⁵ Weber, A. N., Michail, M., Thompson, A., & Fiedorowicz, J. G. (2017). Psychiatric emergencies: Assessing and managing suicidal ideation. *Medical Clinics of North America*, 101(3), 553-571. <https://doi:10.1016/j.mcna.2016.12.006>.
- ³¹⁶ Visit <http://cssrs.columbia.edu/> for more information on the Columbia-Suicide Severity Rating Scale.
- ³¹⁷ Ibid.
- ³¹⁸ Johnson, J. G., Harris, E. S., Spitzer, R. L., & Williams, J. B. (2002). The patient health questionnaire for adolescents: validation of an instrument for the assessment of mental disorders among adolescent primary care patients. *Journal of Adolescent Health*, 30(3), 196-204.
- ³¹⁹ Horowitz, L. M., Bridge, J. A., Teach, S. J., et al. (2012). Ask Suicide-Screening Questions (ASQ): A Brief Instrument for the Pediatric Emergency Department. *Archives of Pediatric and Adolescent Medicine*, 166(12), 1170–1176. <https://doi:10.1001/archpediatrics.2012.1276>.
- ³²⁰ Ibid.
- ³²¹ Stanley, B., & Brown, G. K. (2012). Safety planning intervention: A brief intervention to mitigate suicide risk. *Cognitive and Behavioral Practice*, 19, 256–264.
- ³²² McMyler, C., & Prymachuk, S. (2008). Do 'no-suicide' contracts work? *Journal of Psychiatric and Mental Health Nursing*, 15(6), 512-522. <https://doi:10.1111/j.1365-2850.2008.01286.x>.
- ³²³ Stanley, B., & Brown, G. K. (2012). Safety planning intervention: A brief intervention to mitigate suicide risk. *Cognitive and Behavioral Practice*, 19, 256-264.
- ³²⁴ Bryan, C. J., Mintz, J., Clemans, T. A., Leeson, B., Burch, T. S., Williams, S. R., Maney, E., & Rudd, M.D. (2017). Effect of crisis response planning vs. contracts for safety on suicide risk in U.S. Army Soldiers: A randomized clinical trial. *Journal of Affective Disorders*, 212, 64-72. <https://doi:10.1016/j.jad.2017.01.028>.
- ³²⁵ Jobes, D. A. (2012). The collaborative assessment and management of suicidality (CAMS): An evolving evidence-based clinical approach to suicidal risk. *Suicide and Life-Threatening Behavior*, 42(6), 640–653. <https://doi.org/10.1111/j.1943-278X.2012.00119.x>
- ³²⁶ McCauley, E., Berk, M. S., Asarnow, J. R., et al. (2018). Efficacy of dialectical behavior therapy for adolescents at high risk for suicide: a randomized clinical trial [published online June 20, 2018]. *JAMA Psychiatry*. <https://doi:10.1001/jamapsychiatry.2018.1109>.
- ³²⁷ Miller, A. L., Rathus, J., & Linehan, M. M. (2007). *Dialectical behavior therapy with suicidal adolescents*. The New York: Guilford Press.
- ³²⁸ Mazza, J. J., Dexter-Mazza, E. T., Miller, A. L., Rathus, J. H., Murphy, H. E., & Linehan, M. M. (2016). *The Guilford practical intervention in the schools series. DBT® skills in schools: Skills training for emotional problem solving for adolescents (DBT STEPS-A)*. New York, NY, US: Guilford Press.
- ³²⁹ Stanley, B., Brown, G., Brent, D. A., Wells, K., Poling, K., Curry, J., Kennard, B. D., Wagner, A., Cwik, M. F., Klomek, A. B., Goldstein, T., Vitiello, B., Barnett, S., Daniel, S., ... Hughes, J. (2009). Cognitive-behavioral therapy for suicide prevention (CBT-SP): treatment model, feasibility, and acceptability. *Journal of the American Academy of Child and Adolescent Psychiatry*, 48(10), 1005-1013.
- ³³⁰ Ibid.
- ³³¹ Ibid.
- ³³² Visit <https://cams-care.com/about-cams/>.

-
- ³³³ Jobes, D. A. (2006). *Managing suicidal risk: a collaborative approach*. New York: Guilford Press.
- ³³⁴ Jobes, D. A., Comtois, K. A., Brenner, L. A., Gutierrez, P. M., O'Connor, S. (2016). Lessons learned from clinical trials of the Collaborative Assessment and Management of Suicidality (CAMS) In: O'Connor RC, .Gordon, J. Platt, S.,(eds.). *International handbook of suicide prevention: research, policy, and practice*. 2. West Sussex: Wiley-Blackwell.
- ³³⁵ Park, A., Gysin-Maillart, A., Müller, T. J., Exadaktylos, A., & Michel, K. (2018). Cost-effectiveness of a brief structured intervention program aimed at preventing repeat suicide attempts among those who previously attempted suicide: A secondary analysis of the ASSIP randomized clinical trial. *JAMA Network Open*, 1(6), e183680.
<https://doi.org/10.1001/jamanetworkopen.2018.3680>.
- ³³⁶ Ibid.
- ³³⁷ Ibid.
- ³³⁸ Mann, J. J., Apter, A., Bertolote, J., Beautrais, A., Currier, D., Haas, A., Hegerl, U., Lonnqvist, J., Malone, K., Marusic, A., Mehlum, L., Patton, G., Phillips, M., Rutz, W., Rhimer, Z., Schmidtke, A., Shaffer, D., Silverman, M., Takahasashi, Y., Varnik, A., Wasserman, D., Wip, P., & Hardin, H. (2005). Suicide prevention strategies: A systematic review. *The Journal of the American Medical Association*, 294, 2064–2074.
- ³³⁹ Agency for Health Care Policy and Research. (1999). *Evidence report on treatment of depression: Newer pharmacotherapies*. Washington, DC: AHCPH Evidence-Based Practice Centers.
- ³⁴⁰ Thies-Flechtner, K., Muller-Oerlinghausen, B., Seibert, W., Walther, A., & Greil, W. (1996). Effect of prophylactic treatment on suicide risk in patients with major affective disorders: data from a randomized prospective trial. *Pharmacopsychiatry*, 29, 103-107.
- ³⁴¹ Bernert, R. A., Hom, M. A., Iwata, N. G., & Joiner, T. E. (2017). Objectively assessed sleep variability as an acute warning sign of suicidal ideation in a longitudinal evaluation of young adults at high suicide risk. *The Journal of Clinical Psychiatry*, 78(6), e678-e687.
<https://doi.org/10.4088/JCP.16m11193>.
- ³⁴² The National Institute of Mental Health (NIMH), The National Institute of Health (NIH). (2012). *NCT01689909: Reducing Suicidal Ideation through Insomnia Treatment*. Augusta University: Author.
- ³⁴³ Weissman, C. R., Blumberger, D. M., Brown, P. E., Isserles, M., Rajji, T. K., Downar, J., Mulsant, B. H., Fitzgerald, P. B., Daskalakis, Z. J. (2018). Bilateral Repetitive Transcranial Magnetic Stimulation decreases suicidal ideation in depression. *The Journal of Clinical Psychiatry*, 79 (3), 17m11692. <https://doi.org/10.4088/JCP.17m11692>.
- ³⁴⁴ Lee, J., Narang, P., Enja, M., & Lippmann, S. (2015). Use of ketamine in acute cases of suicidality. *Innovations in clinical neuroscience*, 12(1-2), 29-31.
- ³⁴⁵ Larkin, G. L., Beautrais, A. L. (2011). A preliminary naturalistic study of low-dose ketamine for depression and suicide ideation in the emergency department. *The International Journal of Neuropsychopharmacology*, 14, 1127–1131.
- ³⁴⁶ Larkin, G. L. & Beautrais, A. L. (2010). Emergency departments are underutilized sites for suicide prevention. *Crisis*, 31(1), 1–6.

-
- ³⁴⁷ National Action Alliance for Suicide Prevention Research Prioritization Task Force. (2014). *A Prioritized Research Agenda for Suicide Prevention: An Action Plan to Save Lives*. Retrieved November 5, 2018 from <http://actionallianceforsuicideprevention.org/task-force/research-prioritization>.
- ³⁴⁸ Miller, I. W., Camargo, C. A., Jr., Arias, S. A., Sullivan, A. F., Allen, M. H., Goldstein, A. B., . . . Boudreaux, E. D. (2017). Suicide prevention in an emergency department population: The ED-SAFE study. *JAMA Psychiatry*, 74(6), 563-570. <https://doi:10.1001/jamapsychiatry.2017.0678>.
- ³⁴⁹ Luxton, D. D., June, J. D., & Comtois, K. A. (2013). Can postdischarge follow-up contacts prevent suicide and suicidal behavior? A review of the evidence. *Crisis*, 34, 32-41. <https://doi:10.1027/0227-5910/a000158>.
- ³⁵⁰ Motto, J. A., & Bostrom, A. G. (2001). A randomized controlled trial of post crisis suicide prevention. *Psychiatric Services*, 52(6), 828-833.
- ³⁵¹ Ibid.
- ³⁵² Ibid.
- ³⁵³ Ibid.
- ³⁵⁴ Richardson, J. S., Mark, T. L., & McKeon, R. (2014). The return on investment of post discharge follow-up calls for suicidal ideation or deliberate self-harm. *Psychiatric Services*, 65, 1012-1019.
- ³⁵⁵ Andrews, G., & Sunderland, M. (2009). Telephone case management reduces both distress and psychiatric hospitalization. *Australian and New Zealand Journal of Psychiatry*, 43, 809-811.
- ³⁵⁶ Cvinar, J.G. (2005). Do suicide survivors suffer social stigma: a review of the literature. *Perspectives in Psychiatric Care*, 41, 14-21.
- ³⁵⁷ Survivors of Suicide Loss Task Force. (2015). *Responding to grief, trauma, and distress after a suicide: U.S. National Guidelines*. Washington, DC: National Action Alliance for Suicide Prevention. Retrieved from <http://www.sprc.org/resources-programs/responding-grief-trauma-and-distress-after-suicide-us-national-guidelines>.
- ³⁵⁸ McDaid, C., Trowman, R., Golder, S., Hawton, K., & Sowden, A. (2008). Interventions for people bereaved through suicide: systematic review. *British Journal of Psychiatry*, 193(6), 438-443. <https://doi:10.1192/bjp.bp.107.040824>.
- ³⁵⁹ Kaslow, N. J., Samples, T. C., Rhodes, M., & Gantt, S. (2011). A family-oriented and culturally sensitive postvention approach with suicide survivors. In J. R. Jordan & J. L. McIntosh (Eds.), *Series in death, dying and bereavement. Grief after suicide: Understanding the consequences and caring for the survivors* (pp. 301-323). New York, NY, US: Routledge/Taylor & Francis Group.
- ³⁶⁰ Ibid.